

Hypergraph Turán numbers of vertex disjoint cycles*

Ran Gu, Xueliang Li, Yongtang Shi[†]

Center for Combinatorics and LPMC-TJKLC

Nankai University, Tianjin 300071, China

guran323@163.com, lxl@nankai.edu.cn, shi@nankai.edu.cn

Abstract

The Turán number of a k -uniform hypergraph H , denoted by $ex_k(n; H)$, is the maximum number of edges in any k -uniform hypergraph F on n vertices which does not contain H as a subgraph. Let $\mathcal{C}_\ell^{(k)}$ denote the family of all k -uniform minimal cycles of length ℓ , $\mathcal{S}(\ell_1, \dots, \ell_r)$ denote the family of hypergraphs consisting of unions of r vertex disjoint minimal cycles of length ℓ_1, \dots, ℓ_r , respectively, and $\mathbb{C}_\ell^{(k)}$ denote a k -uniform linear cycle of length ℓ . We determine precisely $ex_k(n; \mathcal{S}(\ell_1, \dots, \ell_r))$ and $ex_k(n; \mathbb{C}_{\ell_1}^{(k)}, \dots, \mathbb{C}_{\ell_r}^{(k)})$ for sufficiently large n . The results extend recent results of Füredi and Jiang [Füredi, Z., Jiang, T. Hypergraph Turán numbers of linear cycles. *J. Combin. Theory Ser. A*, 123(1): 252–270 (2014)], in which the Turán numbers for single k -uniform minimal cycles and linear cycles are determined.

Keywords: Turán number; cycles; extremal hypergraphs

AMS Subject Classification (2010): 05D05, 05C35, 05C65

1 Introduction

In this paper, we employ standard terminology and notation from hypergraph theory (see e.g., [1]). A *hypergraph* is a pair $H = (V, E)$ consisting of a set V of vertices and a set $E \subseteq \mathcal{P}(V)$ of edges. If every edge contains exactly k vertices, then

*Supported by the National Natural Science Foundation of China (No. 11371205, 11371021), 973 program No.2013CB834204, and PCSIRT.

[†]Corresponding author.

H is a k -uniform hypergraph. A graph homomorphism f from a graph $G = (V, E)$ to a graph $G' = (V', E')$ is a mapping $f: V \rightarrow V'$ from the vertex set of G to the vertex set of G' such that $uv \in E$ implies $f(u)f(v) \in E'$. For two hypergraphs G and H , we write $G \subseteq H$ if there is an injective homomorphism from G into H . We use $G \cup H$ to denote the disjoint union of (hyper)graphs G and H . By disjoint, we will always mean vertex disjoint. A *Berge path* of length ℓ is a family of distinct sets $\{F_1, \dots, F_\ell\}$ and $\ell + 1$ distinct vertices $v_1, \dots, v_{\ell+1}$ such that for each $i = 1, 2, \dots, \ell$, F_i contains v_i and v_{i+1} . Let $\mathcal{B}_\ell^{(k)}$ denote the family of k -uniform Berge paths of length ℓ . A *linear path* of length ℓ is a family of sets $\{F_1, \dots, F_\ell\}$ such that $|F_i \cap F_{i+1}| = 1$ for each i and $F_i \cap F_j = \emptyset$ whenever $|i - j| > 1$. Let $\mathbb{P}_\ell^{(k)}$ denote the k -uniform linear path of length ℓ . It is unique up to isomorphisms. A k -uniform *Berge cycle* of length ℓ is a cyclic list of distinct k -sets A_1, \dots, A_ℓ and ℓ distinct vertices v_1, \dots, v_ℓ such that for each $i = 1, 2, \dots, \ell$, A_i contains v_i and v_{i+1} (where $v_{\ell+1} = v_1$). A k -uniform *minimal cycle* of length ℓ is a cyclic list of k -sets A_1, \dots, A_ℓ such that consecutive sets intersect in at least one element and nonconsecutive sets are disjoint. Denote the family of all k -uniform minimal cycles of length ℓ by $\mathcal{C}_\ell^{(k)}$. A k -uniform *linear cycle* of length ℓ , denoted by $\mathbb{C}_\ell^{(k)}$, is a cyclic list of k -sets A_1, \dots, A_ℓ such that consecutive sets intersect in exactly one element and nonconsecutive sets are disjoint.

The *Turán number*, or extremal number, of a k -uniform hypergraph H , denoted by $ex_k(n; H)$, is the maximum number of edges in any k -uniform hypergraph F on n vertices which does not contain H as a subgraph. This is a natural generalization of the classic Turán number for 2-uniform graphs; we restrict ourselves to the case of k -uniform hypergraphs. Let $ex_k(n; F_1, F_2, \dots, F_r)$ denote the *k -uniform hypergraph Turán Number* of a list of k -uniform hypergraphs F_1, F_2, \dots, F_r , i.e., $ex_k(n; F_1, F_2, \dots, F_r) = ex_k(n; F_1 \cup F_2 \cup \dots \cup F_r)$.

For the family of k -uniform Berge paths of length ℓ , Györi, Katona and Lemons [5] determined $ex_k(n; \mathcal{B}_\ell^{(k)})$ exactly for infinitely many n . In [2], Füredi, Jiang and Seiver established the following results.

Theorem 1 ([2]) *Let k, t be positive integers, where $k \geq 3$. For sufficiently large n , we have*

$$ex_k\left(n; \mathbb{P}_{2t+1}^{(k)}\right) = \binom{n-1}{k-1} + \binom{n-2}{k-1} + \dots + \binom{n-t}{k-1}.$$

The only extremal family consists of all the k -sets in $[n]$ that meet some fixed set S of t vertices. Also,

$$ex_k\left(n; \mathbb{P}_{2t+2}^{(k)}\right) = \binom{n-1}{k-1} + \binom{n-2}{k-1} + \dots + \binom{n-t}{k-1} + \binom{n-t-2}{k-2}.$$

The only extremal family consists of all the k -sets in $[n]$ that meet some fixed set S of t vertices plus all the k -sets in $[n] \setminus S$ that contain some two fixed elements.

For more results we refer to [2, 6].

For the minimal and linear cycles, Füredi and Jiang [3] determined the extremal numbers when the forbidden hypergraph is a single minimal cycle or a single linear cycle. This confirms, in a stronger form, a conjecture of Mubayi and Verstraëte [6] for $k \geq 5$ and adds to the limited list of hypergraphs whose Turán numbers have been known either exactly or asymptotically. Their main results are as follows.

Theorem 2 ([3]) *Let t be a positive integer, $k \geq 4$. For sufficiently large n , we have $ex_k(n; \mathcal{C}_{2t+1}^{(k)}) = \binom{n}{k} - \binom{n-t}{k}$, and $ex_k(n; \mathcal{C}_{2t+2}^{(k)}) = \binom{n}{k} - \binom{n-t}{k} + 1$. For $\mathcal{C}_{2t+1}^{(k)}$, the only extremal family consists of all the k -sets in $[n]$ that meet some fixed t -set S . For $\mathcal{C}_{2t+2}^{(k)}$, the only extremal family consists of all the k -sets in $[n]$ that intersect some fixed t -set S plus one additional k -set outside S .*

Theorem 3 ([3]) *Let t be a positive integer, $k \geq 5$. For sufficiently large n , we have $ex_k(n; \mathcal{C}_{2t+1}^{(k)}) = \binom{n}{k} - \binom{n-t}{k}$, and $ex_k(n; \mathcal{C}_{2t+2}^{(k)}) = \binom{n}{k} - \binom{n-t}{k} + \binom{n-t-2}{k-2}$. For $\mathcal{C}_{2t+1}^{(k)}$, the only extremal family consists of all the k -sets in $[n]$ that meet some fixed t -set S . For $\mathcal{C}_{2t+2}^{(k)}$, the only extremal family consists of all the k -sets in $[n]$ that intersect some fixed t -set S plus all the k -sets in $[n] \setminus S$ that contain some two fixed elements.*

From definition, two k -uniform minimal cycles of the same length may not be isomorphic. Hence, we define the following family of hypergraphs, where every member consists of r vertex disjoint cycles:

$$\mathcal{S}(\ell_1, \dots, \ell_r) = \{C_1 \cup \dots \cup C_r : C_i \in \mathcal{C}_{\ell_i}^{(k)} \text{ for } i \in [r]\}.$$

Apart from the results above, we will need the following result, due to Keevash, Mubayi and Wilson [4].

Theorem 4 ([4]) *Let H be a k -uniform hypergraph on n vertices with no two edges intersecting in exactly one vertex, where $k \geq 3$. Then $|E(H)| \leq \binom{n}{k-2}$.*

Based on earlier work of Füredi and Jiang [3], in this paper we will determine precisely the exact Turán numbers when the forbidden hypergraphs are r vertex disjoint minimal cycles or r vertex disjoint linear cycles. Our main results are as follows.

Theorem 5 *Let integers $k \geq 4$, $r \geq 1$, $\ell_1, \dots, \ell_r \geq 3$, $t = \sum_{i=1}^r \lfloor \frac{\ell_i+1}{2} \rfloor - 1$, and $I = 1$ if all the ℓ_1, \dots, ℓ_r are even, and $I = 0$ otherwise. For sufficiently large n ,*

$$ex_k(n; \mathcal{S}(\ell_1, \dots, \ell_r)) = \binom{n}{k} - \binom{n-t}{k} + I.$$

Theorem 6 *Let integers $k \geq 5$, $r \geq 1$, $\ell_1, \dots, \ell_r \geq 3$, $t = \sum_{i=1}^r \lfloor \frac{\ell_i+1}{2} \rfloor - 1$, and $J = \binom{n-t-2}{k-2}$ if all the ℓ_1, \dots, ℓ_r are even, and $J = 0$ otherwise. For sufficiently large n ,*

$$ex_k(n; \mathbb{C}_{\ell_1}^{(k)}, \dots, \mathbb{C}_{\ell_r}^{(k)}) = \binom{n}{k} - \binom{n-t}{k} + J.$$

Sometimes, we allow the hypergraph to contain less than r minimal or linear cycles, consider the Turán number in such cases, we have the following two corollaries. We use notation $r \cdot F$ to denote r vertex disjoint copies of a hypergraph F . Let $\ell_1 = \dots = \ell_r = \ell$, we can immediately get the following two corollaries from Theorems 5 and 6.

Corollary 1 *Let integers $k \geq 4$, $r \geq 1$, $\ell \geq 3$, $t = r \lfloor \frac{\ell+1}{2} \rfloor - 1$, and $I = 1$ if ℓ is even, and $I = 0$ if ℓ is odd. For sufficiently large n ,*

$$ex_k(n; r \cdot \mathbb{C}_{\ell}^{(k)}) = \binom{n}{k} - \binom{n-t}{k} + I.$$

Corollary 2 *Let integers $k \geq 5$, $r \geq 1$, $\ell \geq 3$, $t = r \lfloor \frac{\ell+1}{2} \rfloor - 1$, and $J = \binom{n-t-2}{k-2}$ if ℓ is even, and $J = 0$ if ℓ is odd. For sufficiently large n ,*

$$ex_k(n; r \cdot \mathbb{C}_{\ell}^{(k)}) = \binom{n}{k} - \binom{n-t}{k} + J.$$

We can see that Theorem 2 and Theorem 3 are special cases of Corollary 1 and Corollary 2 (when $r = 1$), respectively. However, the methods we used in the proofs of Theorem 5 and Theorem 6 are quite different from Theorem 2 and Theorem 3.

2 Proof of Theorem 5

For convenience, we define $f(n, k, \{\ell_1, \dots, \ell_r\}) = \binom{n}{k} - \binom{n-t}{k} + I$. Note that the hypergraph on n vertices that has every edge incident to some fixed t -set S , along with one additional edge disjoint from S when all of ℓ_1, \dots, ℓ_r are even, has exactly $f(n, k, \{\ell_1, \dots, \ell_r\})$ edges and does not contain a copy of any member of $\mathcal{S}(\ell_1, \dots, \ell_r)$.

Thus, to prove Theorem 5, it suffices to show that $ex_k(n; \mathcal{S}(\ell_1, \dots, \ell_r)) \leq \binom{n}{k} - \binom{n-t}{k} + I$, i.e., any hypergraph on n vertices with more than $f(n, k, \{\ell_1, \dots, \ell_r\})$ edges must contain a member of $\mathcal{S}(\ell_1, \dots, \ell_r)$. We use induction on r . From Theorem 2, the case $r = 1$ has been proved. Assume that $r \geq 2$, and Theorem 5 holds for smaller r .

Let H be a hypergraph on n vertices with m edges and $m > f(n, k, \{\ell_1, \dots, \ell_r\})$. Since $f(n, k, \{\ell_1, \dots, \ell_r\}) > f(n, k, \ell_1)$ for sufficiently large n , there exists at least one k -uniform minimal ℓ_1 -cycle in H . Take one of them, denote its vertex set by C , so $\ell_1 \leq |C| \leq (k-1)\ell_1$. We have $|E(H \setminus C)| \leq f(n - |C|, k, \{\ell_2, \dots, \ell_r\})$, since otherwise, by induction hypothesis, we can find vertex disjoint copies of $\mathcal{C}_{\ell_2}^{(k)} \cup \dots \cup \mathcal{C}_{\ell_r}^{(k)}$ in H , plus the minimal ℓ_1 -cycle on C , and then there is a copy of a member of $\mathcal{S}(\ell_1, \dots, \ell_r)$ in H already.

Let m_C denote the number of edges in H incident to vertices in C . Then,

$$m_C \geq m - f(n - |C|, k, \{\ell_2, \dots, \ell_r\}) \quad (1)$$

$$\geq f(n, k, \{\ell_1, \dots, \ell_r\}) - f(n - \ell_1, k, \{\ell_2, \dots, \ell_r\}) \quad (2)$$

$$= \frac{\lfloor \frac{\ell_1+1}{2} \rfloor}{(k-1)!} n^{k-1} + O(n^{k-2}). \quad (3)$$

We call an edge in H a *terminal edge* if it contains exactly one vertex in C . Let T denote the set of all terminal edges in H . For every $(k-1)$ -set R in $V(H) \setminus C$, define

$$T_R = \{E \in T : R \subseteq E\}.$$

According to the size of each set T_R , we divide all the $(k-1)$ -sets in $V(H) \setminus C$ into two sets, such that

$$X = \{R \subseteq V(H) \setminus C \text{ and } |R| = k-1 : |T_R| \leq \left\lfloor \frac{\ell_1+1}{2} \right\rfloor - 1\}$$

$$Y = \{R \subseteq V(H) \setminus C \text{ and } |R| = k-1 : |T_R| \geq \left\lfloor \frac{\ell_1+1}{2} \right\rfloor\}.$$

It is not difficult to give an upper bound of m_C in terms of $|X|$ and $|Y|$ as follows:

$$\begin{aligned} m_C &\leq \binom{|C|}{2} \binom{n-2}{k-2} + |X| \left(\left\lfloor \frac{\ell_1+1}{2} \right\rfloor - 1 \right) + |Y| \cdot |C| \\ &\leq \binom{|C|}{2} \binom{n-2}{k-2} + \binom{n}{k-1} \left(\left\lfloor \frac{\ell_1+1}{2} \right\rfloor - 1 \right) + |Y| \cdot \ell_1 (k-1). \end{aligned}$$

Combine with (3), we have

$$|Y| \geq \frac{n^{k-1}}{(k-1)\ell_1(k-1)!} + O(n^{k-2}). \quad (4)$$

For any $(k-1)$ -set $R \in Y$, there are at least $\lfloor \frac{\ell_1+1}{2} \rfloor$ vertices in C that can form terminal edges with R . We choose exactly $\lfloor \frac{\ell_1+1}{2} \rfloor$ of them, and call the vertex set of these $\lfloor \frac{\ell_1+1}{2} \rfloor$ vertices *terminal set* relative to R . Since the number of $\lfloor \frac{\ell_1+1}{2} \rfloor$ -sets in C is at most $\binom{|C|}{\lfloor \frac{\ell_1+1}{2} \rfloor}$, we can get that some elements in Y may have the same terminal set. And it is easy to derive that the number of $(k-1)$ -sets in Y with the same terminal set is at least

$$\frac{n^{k-1}}{(k-1)\ell_1(k-1)!} \binom{|C|}{\lfloor \frac{\ell_1+1}{2} \rfloor}^{-1} + O(n^{k-2}) \geq \frac{n^{k-1}}{(k-1)\ell_1(k-1)!} \left(\binom{(k-1)\ell_1}{\lfloor \frac{\ell_1+1}{2} \rfloor} \right)^{-1} + O(n^{k-2}).$$

Choose one terminal set U in C , such that there are at least $\frac{n^{k-1}}{(k-1)\ell_1(k-1)!} \left(\binom{(k-1)\ell_1}{\lfloor \frac{\ell_1+1}{2} \rfloor} \right)^{-1} + O(n^{k-2})$ $(k-1)$ -sets in $V(H) \setminus C$, every such $(k-1)$ -set can form a terminal edge with every vertex in U . Let R_U be the set of all the common $(k-1)$ -sets associated with U in $V(H) \setminus C$. Then we have

$$|R_U| \geq \frac{n^{k-1}}{(k-1)\ell_1(k-1)!} \left(\binom{(k-1)\ell_1}{\lfloor \frac{\ell_1+1}{2} \rfloor} \right)^{-1} + O(n^{k-2}). \quad (5)$$

Let m_U denote the number of edges incident to vertices in U . Then,

$$m_U \leq \left\lfloor \frac{\ell_1+1}{2} \right\rfloor \binom{n - \lfloor \frac{\ell_1+1}{2} \rfloor}{k-1} + m',$$

where m' is the number of edges which contain at least two vertices in U . With some calculations, we have

$$\begin{aligned} &f(n, k, \{\ell_1, \dots, \ell_r\}) - f\left(n - \left\lfloor \frac{\ell_1+1}{2} \right\rfloor, k, \{\ell_2, \dots, \ell_r\}\right) - m_U \\ &= \binom{n-1}{k-1} + \binom{n-2}{k-1} + \dots + \binom{n - \lfloor \frac{\ell_1+1}{2} \rfloor}{k-1} - m_U \\ &\geq \left[\binom{n-1}{k-1} - \binom{n - \lfloor \frac{\ell_1+1}{2} \rfloor}{k-1} \right] + \left[\binom{n-2}{k-1} - \binom{n - \lfloor \frac{\ell_1+1}{2} \rfloor}{k-1} \right] \\ &\quad + \dots + \left[\binom{n - \lfloor \frac{\ell_1+1}{2} \rfloor + 1}{k-1} - \binom{n - \lfloor \frac{\ell_1+1}{2} \rfloor}{k-1} \right] - m'. \end{aligned}$$

It is not difficult to deduce that the last expression is nonnegative (consider the combinatorial meaning of that expression). Hence, we can derive

$$\begin{aligned} E(H \setminus U) &= m - m_U > f(n, k, \{\ell_1, \dots, \ell_r\}) - m_U \\ &\geq f\left(n - \left\lfloor \frac{\ell_1 + 1}{2} \right\rfloor, k, \{\ell_2, \dots, \ell_r\}\right). \end{aligned}$$

Thus by the induction hypothesis, there exists a member of $\mathcal{S}(\ell_2, \dots, \ell_r)$ with vertex set W in $V(H) \setminus U$. Also, we have

$$|W| \leq (k-1) \sum_{i=2}^r \ell_i. \quad (6)$$

Now we focus on finding a k -uniform minimal ℓ_1 -cycle disjoint from W . Considering the $(k-1)$ -uniform hypergraph H_0 with vertex set $V(H) \setminus U$ and edge set R_U , we will prove the following claim:

Claim 1 *There are $\lfloor \frac{\ell_1}{2} \rfloor$ pairs of $(k-1)$ -edges in H_0 , say $\{a_i, b_i\}$, $i = 1, \dots, \lfloor \frac{\ell_1}{2} \rfloor$, such that for every i , a_i and b_i have exactly one common vertex, and for any $j \neq i$, $\{a_i, b_i\}$ and $\{a_j, b_j\}$ are vertex disjoint, moreover, all these $(k-1)$ -edges disjoint from W .*

Proof. The number of $(k-1)$ -edges incident with some vertices in W is at most $|W| \cdot \binom{n-1}{k-2}$. With the aid of (5) and (6), in R_U the number of $(k-1)$ -edges disjoint from W is at least

$$\frac{n^{k-1}}{(k-1)\ell_1(k-1)!} \left(\binom{(k-1)\ell_1}{\lfloor \frac{\ell_1+1}{2} \rfloor} \right)^{-1} + O(n^{k-2}) - (k-1) \sum_{i=2}^r \ell_i \binom{n-1}{k-2} > \binom{n - \lfloor \frac{\ell_1+1}{2} \rfloor}{k-2}.$$

By Theorem 4, we can find a pair $\{a_1, b_1\}$ of $(k-1)$ -edges with exactly one common vertex. Let $p = \lfloor \frac{\ell_1}{2} \rfloor (2k-3)$. Since $\frac{n^{k-1}}{(k-1)\ell_1(k-1)!} \left(\binom{(k-1)\ell_1}{\lfloor \frac{\ell_1+1}{2} \rfloor} \right)^{-1} + O(n^{k-2}) - (k-1) \sum_{i=2}^r \ell_i \binom{n-1}{k-2} - p \binom{n-1}{k-2} > \binom{n - \lfloor \frac{\ell_1+1}{2} \rfloor}{k-2}$, we can repeat the argument above to find $\{a_2, b_2\}, \dots, \{a_{\lfloor \frac{\ell_1}{2} \rfloor}, b_{\lfloor \frac{\ell_1}{2} \rfloor}\}$ satisfying the properties described in Claim 1. \square

Let $U = \{u_1, \dots, u_{\lfloor \frac{\ell_1+1}{2} \rfloor}\}$. To form the required minimal ℓ_1 -cycle, we need to consider the following two cases:

Case 1. ℓ_1 is even.

Find $\frac{\ell_1}{2}$ pairs of $(k-1)$ -edges in H_0 as described in Claim 1, still denote them by $\{a_i, b_i\}$, $i = 1, \dots, \frac{\ell_1}{2}$. Construct a k -uniform minimal ℓ_1 -cycle in H with edges:

$$a_1 \cup \{u_1\}, b_1 \cup \{u_2\}, a_2 \cup \{u_2\}, \dots, b_{\frac{\ell_1}{2}-1} \cup \{u_{\frac{\ell_1}{2}}\}, a_{\frac{\ell_1}{2}} \cup \{u_{\frac{\ell_1}{2}}\}, b_{\frac{\ell_1}{2}} \cup \{u_1\}.$$

Case 2. ℓ_1 is odd.

Find $\frac{\ell_1-3}{2}$ pairs of $(k-1)$ -edges in H_0 as described in Claim 1. Similar to the proof of Claim 1. Let Q be the union of W and the set of vertices in all these $\frac{\ell_1-3}{2}$ pairs of $(k-1)$ -edges. Hence, $|Q| = \frac{\ell_1-3}{2}(2k-3) + |W|$. By Theorem 1, $ex_{k-1}\left(n - \lfloor \frac{\ell_1+1}{2} \rfloor; \mathbb{P}_3^{(k-1)}\right) = \frac{1}{(k-2)!}n^{k-2} + O(n^{k-3})$, for sufficiently large n . In H_0 , the number of $(k-1)$ -edges disjoint from Q is at least $\frac{n^{k-1}}{(k-1)\ell_1(k-1)!} \binom{(k-1)\ell_1}{\lfloor \frac{\ell_1+1}{2} \rfloor}^{-1} + O(n^{k-2}) - |Q| \binom{n-1}{k-2} > \frac{1}{(k-2)!}n^{k-2} + O(n^{k-3})$. That implies that in H_0 we can find a $\mathbb{P}_3^{(k-1)}$ in the remaining $(k-1)$ -edges disjoint from Q . Let x, y, z be the three consecutive $(k-1)$ -edges in $\mathbb{P}_3^{(k-1)}$. Then, in H we can form a k -uniform minimal ℓ_1 -cycle with edges:

$$\begin{aligned} a_1 \cup \{u_1\}, b_1 \cup \{u_2\}, a_2 \cup \{u_2\}, \dots, a_{\frac{\ell_1-3}{2}} \cup \{u_{\frac{\ell_1-3}{2}}\}, \\ b_{\frac{\ell_1-3}{2}} \cup \{u_{\frac{\ell_1-1}{2}}\}, x \cup \{u_{\frac{\ell_1-1}{2}}\}, y \cup \{u_{\frac{\ell_1+1}{2}}\}, z \cup \{u_1\}. \end{aligned}$$

Moreover, it is easy to see that this k -uniform minimal ℓ_1 -cycle is not only minimal, but also linear, no matter when ℓ_1 is even or odd. Thus, we have constructed r disjoint k -uniform minimal cycles. So, the hypergraph which contains no member of $\mathcal{S}(\ell_1, \dots, \ell_r)$ can not have more than $f(n, k, \{\ell_1, \dots, \ell_r\})$ edges. The proof is thus complete. \blacksquare

3 Proof of Theorem 6

Let $g(n, k, \{\ell_1, \dots, \ell_r\}) = \binom{n}{k} - \binom{n-t}{k} + J$. Firstly, we point out that the hypergraph on n vertices that has every edge incident to some fixed t -set S , along with all the k -edges disjoint from S containing some two fixed elements not in S when all of ℓ_1, \dots, ℓ_r are even, has exactly $g(n, k, \{\ell_1, \dots, \ell_r\})$ edges and does not contain a copy of any member of $\mathbb{C}_{\ell_1}^{(k)} \cup \dots \cup \mathbb{C}_{\ell_r}^{(k)}$.

Hence, it suffices to show that $ex_k\left(n; \mathbb{C}_{\ell_1}^{(k)}, \dots, \mathbb{C}_{\ell_r}^{(k)}\right) \leq g(n, k, \{\ell_1, \dots, \ell_r\})$. We proceed by induction on r again since the case $r = 1$ is provided by Theorem 3. Let H be a hypergraph on n vertices with $m > g(n, k, \{\ell_1, \dots, \ell_r\})$ edges. If one of ℓ_1, \dots, ℓ_r is even, rearrange the sequence to make sure ℓ_1 is even.

As in the proof of Theorem 5, since $g(n, k, \{\ell_1, \dots, \ell_r\}) > g(n, k, \ell_1)$ for sufficiently large n , there exists at least one k -uniform linear ℓ_1 -cycle in H . Take one of them, denote its vertex set by C . Similarly, we have $|E(H \setminus C)| \leq g(n - |C|, k, \{\ell_2, \dots, \ell_r\})$. Still let m_C denote the number of edges in H incident to some vertices in C . With

some calculations, we can get

$$m_C \geq \frac{\lfloor \frac{\ell_1+1}{2} \rfloor}{(k-1)!} n^{k-1} + O(n^{k-2}).$$

Again we define terminal edges, T_R, X, Y as before, we then can find the $\lfloor \frac{\ell_1+1}{2} \rfloor$ -set U , too. Then by induction hypothesis, we can find a copy of $\mathbb{C}_{\ell_2}^{(k)} \cup \dots \cup \mathbb{C}_{\ell_r}^{(k)}$ on vertex set W in $V(H) \setminus U$. Now we focus on finding a k -uniform linear ℓ_1 -cycle disjoint from W . Again considering the $(k-1)$ -uniform hypergraph H_0 with vertex set $V(H) \setminus U$ and edge set R_U , it is easy to see that the Claim 1 still holds. Thus, like Theorem 5, we have the terminal set $U = \{u_1, \dots, u_{\lfloor \frac{\ell_1+1}{2} \rfloor}\}$. To form the required linear ℓ_1 -cycle, we also need to consider the following two cases:

Case 1. ℓ_1 is even.

Find $\frac{\ell_1}{2}$ pairs of $(k-1)$ -edges in H_0 as described in Claim 1, still denote them by $\{a_i, b_i\}$, $i = 1, \dots, \frac{\ell_1}{2}$. Construct a k -uniform linear ℓ_1 -cycle in H with edges:

$$a_1 \cup \{u_1\}, b_1 \cup \{u_2\}, a_2 \cup \{u_2\}, \dots, b_{\frac{\ell_1}{2}-1} \cup \{u_{\frac{\ell_1}{2}}\}, a_{\frac{\ell_1}{2}} \cup \{u_{\frac{\ell_1}{2}}\}, b_{\frac{\ell_1}{2}} \cup \{u_1\}.$$

Case 2. ℓ_1 is odd.

Find $\frac{\ell_1-3}{2}$ pairs of $(k-1)$ -edges in H_0 as described in Claim 1. Similar to the proof of Claim 1. Let Q be the union of W and the set of vertices in all these $\frac{\ell_1-3}{2}$ pairs of $(k-1)$ -edges. Hence, $|Q| = \frac{\ell_1-3}{2}(2k-3) + |W|$. By Theorem 1, $ex_{k-1}\left(n - \lfloor \frac{\ell_1+1}{2} \rfloor; \mathbb{P}_3^{(k-1)}\right) = \frac{1}{(k-2)!} n^{k-2} + O(n^{k-3})$, for sufficiently large n . In H_0 , the number of $(k-1)$ -edges disjoint from Q is at least $\frac{n^{k-1}}{(k-1)\ell_1(k-1)!} \binom{(k-1)\ell_1}{\lfloor \frac{\ell_1+1}{2} \rfloor}^{-1} + O(n^{k-2}) - |Q| \binom{n-1}{k-2} > \frac{1}{(k-2)!} n^{k-2} + O(n^{k-3})$. That implies that in H_0 we can find a $\mathbb{P}_3^{(k-1)}$ in the remaining $(k-1)$ -edges disjoint from Q . Let x, y, z be the three consecutive $(k-1)$ -edges in $\mathbb{P}_3^{(k-1)}$. Then, in H we can form a k -uniform linear ℓ_1 -cycle with edges:

$$\begin{aligned} & a_1 \cup \{u_1\}, b_1 \cup \{u_2\}, a_2 \cup \{u_2\}, \dots, a_{\frac{\ell_1-3}{2}} \cup \{u_{\frac{\ell_1-3}{2}}\}, \\ & b_{\frac{\ell_1-3}{2}} \cup \{u_{\frac{\ell_1-1}{2}}\}, x \cup \{u_{\frac{\ell_1-1}{2}}\}, y \cup \{u_{\frac{\ell_1+1}{2}}\}, z \cup \{u_1\}. \end{aligned}$$

Since we construct this k -uniform linear ℓ_1 -cycle avoiding the vertices in W , we know that the hypergraph containing no $\mathbb{C}_{\ell_1}^{(k)} \cup \dots \cup \mathbb{C}_{\ell_r}^{(k)}$ can not have more than $g(n, k, \{\ell_1, \dots, \ell_r\})$ edges. The proof is then complete. \blacksquare

References

- [1] Berge, C. Graphs and Hypergraphs. North-Holland Publishing, North-Holland, Amsterdam, 1989.

- [2] Füredi, Z., Jiang, T., Seiver, R. Exact solution of the hypergraph Turán problem for k -uniform linear paths. *Combinatorica*, 34(3): 299-322 (2014)
- [3] Füredi, Z., Jiang, T. Hypergraph Turán numbers of linear cycles. *J. Combin. Theory Ser. A*, 123(1): 252–270 (2014)
- [4] Keevash, P., Mubayi, D., Wilson, R. Set systems with no singleton intersection. *SIAM J. Discrete Math.*, 20: 1031–1041 (2006)
- [5] Györi, E., Katona, G., Lemons, N. Hypergraph extensions of the Erdős-Gallai theorem. *Electronic Notes in Discrete Math.*, 36: 655–662 (2010)
- [6] Mubayi, D., Verstraëte, J. Minimal paths and cycles in set systems. *European J. Combin.*, 28: 1681–1693 (2007)