

Extremal Theta-free planar graphs

Yongxin Lan¹, Yongtang Shi^{2,*}, Zi-Xia Song³

¹*School of Mathematical Sciences, Tianjin Polytechnic University, Tianjin, 300387, China*

²*Center for Combinatorics and LPMC, Nankai University, Tianjin 300071, China*

³*Department of Mathematics, University of Central Florida, Orlando, FL 32816, USA*

Abstract

Given a family \mathcal{F} , a graph is \mathcal{F} -free if it does not contain any graph in \mathcal{F} as a subgraph. We continue to study the topic of “extremal” planar graphs initiated by Dowden [J. Graph Theory 83 (2016) 213–230], that is, how many edges can an \mathcal{F} -free planar graph on n vertices have? We define $ex_{\mathcal{P}}(n, \mathcal{F})$ to be the maximum number of edges in an \mathcal{F} -free planar graph on n vertices. Dowden obtained the tight bounds $ex_{\mathcal{P}}(n, C_4) \leq 15(n-2)/7$ for all $n \geq 4$ and $ex_{\mathcal{P}}(n, C_5) \leq (12n-33)/5$ for all $n \geq 11$. In this paper, we continue to promote the idea of determining $ex_{\mathcal{P}}(n, \mathcal{F})$ for certain classes \mathcal{F} . Let Θ_k denote the family of Theta graphs on $k \geq 4$ vertices, that is, graphs obtained from a cycle C_k by adding an additional edge joining two non-consecutive vertices. The study of $ex_{\mathcal{P}}(n, \Theta_4)$ was suggested by Dowden. We show that $ex_{\mathcal{P}}(n, \Theta_4) \leq 12(n-2)/5$ for all $n \geq 4$, $ex_{\mathcal{P}}(n, \Theta_5) \leq 5(n-2)/2$ for all $n \geq 5$, and then demonstrate that these bounds are tight, in the sense that there are infinitely many values of n for which they are attained exactly. We also prove that $ex_{\mathcal{P}}(n, C_6) \leq ex_{\mathcal{P}}(n, \Theta_6) \leq 18(n-2)/7$ for all $n \geq 6$.

AMS Classification: 05C10; 05C35.

Keywords: Turán number; extremal graph; planar graph

1 Introduction

All graphs considered in this paper are finite and simple. We use P_k and C_k to denote the path and cycle on k vertices, respectively. Let \mathcal{F} be a family of graphs. A graph is \mathcal{F} -free if it does not contain any graph in \mathcal{F} as a subgraph. When $\mathcal{F} = \{F\}$ we write F -free. One of the best known results in extremal graph theory is Turán’s Theorem [12], which gives the maximum number of edges that a K_k -free graph on n vertices can have. The celebrated Erdős-Stone Theorem [4] then extends this to the case when K_k is replaced by an arbitrary graph H , showing that the maximum number of edges possible is $(1 + o(1)) \left(\frac{\chi(H)-2}{\chi(H)-1} \right) n$, where $\chi(H)$ denotes the chromatic number of H . This latter result has been called the “fundamental theorem of extremal graph theory” [1]. Turán-type problems when host graphs are hypergraphs are notoriously difficult. A large quantity

*E-mail addresses: yxlan0@126.com (Y. Lan); shi@nankai.edu.cn (Y. Shi); Zixia.Song@ucf.edu (Z-X. Song)

of work in this area has been carried out in determining the maximum number of edges in a k -uniform hypergraph on n vertices without containing k -uniform linear paths and cycles (see, for example, [6, 7, 10]). Surveys on Turán-type problems of graphs and hypergraphs can be found in [5] and [9].

Recently, Dowden [3] initiated the study of Turán-type problems when host graphs are planar graphs, i.e., how many edges can an \mathcal{F} -free planar graph on n vertices have? The *planar Turán number of \mathcal{F}* , denoted $ex_{\mathcal{P}}(n, \mathcal{F})$, is the maximum number of edges in an \mathcal{F} -free planar graph on n vertices. When $\mathcal{F} = \{F\}$ we write $ex_{\mathcal{P}}(n, F)$. Dowden [3] observed that it is straightforward to determine the exact values of $ex_{\mathcal{P}}(n, H)$ when H is a complete graph or non-planar graph; he also obtained the tight bounds $ex_{\mathcal{P}}(n, C_4) \leq 15(n - 2)/7$ for all $n \geq 4$ and $ex_{\mathcal{P}}(n, C_5) \leq (12n - 33)/5$ for all $n \geq 11$. Recently, Lan, Shi and Song observed in [11] that planar Turán numbers are closely related to planar anti-Ramsey numbers. The *planar anti-Ramsey number of \mathcal{F}* , denoted $ar_{\mathcal{P}}(n, \mathcal{F})$, is the maximum number of colors in an edge-coloring of a plane triangulation T (which is not \mathcal{F} -free) on n vertices such that T contains no rainbow copy of any $F \in \mathcal{F}$. When $\mathcal{F} = \{F\}$ we write $ar_{\mathcal{P}}(n, F)$. The study of planar anti-Ramsey numbers was initiated by Horňák, Jendrol', Schiermeyer and Soták [8] (under the name of rainbow numbers). The following result is observed in [11].

Proposition 1.1 ([11]) *Given a planar graph H and a positive integer $n \geq |H|$,*

$$1 + ex_{\mathcal{P}}(n, \mathcal{H}) \leq ar_{\mathcal{P}}(n, H) \leq ex_{\mathcal{P}}(n, H),$$

where $\mathcal{H} = \{H - e : e \in E(H)\}$.

In this paper, we continue to promote the idea of determining $ex_{\mathcal{P}}(n, \mathcal{F})$ for certain classes \mathcal{F} . This paper focuses on the family of Theta graphs, where a graph on at least 4 vertices is a *Theta graph* if it can be obtained from a cycle by adding an additional edge joining two non-consecutive vertices. For integer $k \geq 4$, let Θ_k be the family of non-isomorphic Theta graphs on k vertices. Note that the only graph in Θ_4 is isomorphic to K_4 minus one edge, and Θ_5 has only one graph. By abusing notation, we also use Θ_4 and Θ_5 to denote the only graph in Θ_4 and Θ_5 , respectively. Note that the study of $ex_{\mathcal{P}}(n, \Theta_4)$ was suggested by Dowden [3]. We need to introduce more notation. For a graph G , we will use $V(G)$ to denote the vertex set, $E(G)$ the edge set, $|G|$ the number of vertices, $e(G)$ the number of edges, $\delta(G)$ the minimum degree and \overline{G} the complement of G . For a vertex $x \in V(G)$, we will use $N_G(x)$ to denote the set of vertices in G which are adjacent to x . We define $d_G(x) = |N_G(x)|$. Given vertex sets $A, B \subseteq V(G)$, the subgraph of G induced on A , denoted $G[A]$, is the graph with vertex set A and edge set $\{xy \in E(G) : x, y \in A\}$. We denote by $B \setminus A$ the set $B - A$ and $G \setminus A$ the subgraph of G induced on $V(G) \setminus A$, respectively. We say that A is *complete to* (resp. *anti-complete to*) B if for every $a \in A$ and every $b \in B$, $ab \in E(G)$ (resp. $ab \notin E(G)$).

If $A = \{a\}$, we simply say a is complete to (resp. anti-complete to) B , and write $B \setminus a$ and $G \setminus a$, respectively. The *join* $G + H$ (resp. *union* $G \cup H$) of two vertex disjoint graphs G and H is the graph having vertex set $V(G) \cup V(H)$ and edge set $E(G) \cup E(H) \cup \{xy \mid x \in V(G), y \in V(H)\}$. (resp. $E(G) \cup E(H)$). For a positive integer t , we use tH to denote disjoint union of t copies of a graph H . Given two isomorphic graphs G and H , we may (with a slight but common abuse of notation) write $G = H$. A graph H is a *spanning subgraph* of a graph G if H is a subgraph of G with $V(H) = V(G)$. For any positive integer k , let $[k] := \{1, 2, \dots, k\}$.

We state and prove our main results in Section 2.

2 Planar Turán number of Theta graphs

In this section, using the method developed in [3], we study planar Turán numbers of Θ_k when $k \in \{4, 5, 6\}$. The study of $ex_{\mathcal{P}}(n, \Theta_4)$ was suggested by Dowden [3]. Our technique relies heavily on Euler's formula. We need to introduce more notation that shall be used in this section only.

An \mathcal{F} -free planar graph G on n vertices with the largest possible number of edges is called *extremal for n and \mathcal{F}* . If $\mathcal{F} = \{F\}$, then we simply say G is extremal for n and F . Given a plane graph G and integers $i, j \geq 3$, an *i -face* in G is a face of size i ; and let: $E_{i,j}$ denote the set of edges in G that each belong to one i -face and one j -face (and belong to two i -faces when $i = j$); E_i denote the set of edges in G that each belong to at least one i -face; and f_i denote the number of i -faces in G . Let $e_{i,j} := |E_{i,j}|$, $e_i := |E_i|$, and $f := \sum_i f_i$. Given three positive integers a, b and c , we use $a \equiv b \pmod{c}$ to denote a and b have the same remainder when divided by c . We will make use of the following observation.

Observation 2.1 *Let G be a plane graph on $n \geq 3$ vertices with $e(G) \geq 2$. For all $i \geq 3$,*

- (a) $e_{i,i} \leq e_i \leq e(G)$,
- (b) $if_i = e_i + e_{i,i}$,
- (c) $\sum_{i \geq 3} e_i - \sum_{3 \leq i < j} e_{i,j} = e(G)$, and
- (d) *every face in G is bounded by a cycle if G is 2-connected.*

We begin with $\mathcal{F} = \Theta_4$ and prove that $ex_{\mathcal{P}}(n, \Theta_4) \leq 12(n-2)/5$ for all $n \geq 4$ and then demonstrate that this bound is tight, in the sense that there are infinitely many values of n for which it is attained exactly.

Theorem 2.2 *$ex_{\mathcal{P}}(n, \Theta_4) \leq 12(n-2)/5$ for all $n \geq 4$, with equality when $n \equiv 12 \pmod{20}$.*

Proof. Let G be a Θ_4 -free plane graph on $n \geq 4$ vertices. We shall proceed the proof by induction on n . The statement is trivially true when $n = 4$ because any Θ_4 -free plane graph on

four vertices has at most four edges. So we may assume that $n \geq 5$. Next assume that there exists a vertex $u \in V(G)$ with $d_G(u) \leq 2$. By the induction hypothesis, $e(G \setminus u) \leq 12(n - 3)/5$ and so $e(G) = e(G \setminus u) + d_G(u) \leq 12(n - 3)/5 + 2 < 12(n - 2)/5$, as desired. So we may assume that $\delta(G) \geq 3$. Then each component of G has at least five vertices because G is Θ_4 -free. By the induction hypothesis, we may further assume that G is connected. Then G has no face of size at most two because G is simple. Hence

$$2e(G) = \sum_{i \geq 3} i f_i \geq 3f_3 + 4 \sum_{i \geq 4} f_i = 3f_3 + 4(f - f_3) = 4f - f_3, \tag{1}$$

which implies that $f \leq (2e(G) + f_3)/4$. Note that $E_{3,3} = \emptyset$ else G would contain Θ_4 as a subgraph, a contradiction. Thus $e_3 = 3f_3$ by Observation 2.1(b). This, together with $e_3 \leq e(G)$ and $f \leq (2e(G) + f_3)/4$, implies that $f \leq 7e(G)/12$. By Euler's formula, $n - 2 = e(G) - f \geq 5e(G)/12$. Hence $e(G) \leq 12(n - 2)/5$, as desired.




Figure 1: Construction of G_k .

From the proof above, we see that equality in $e(G) \leq 12(n - 2)/5$ is achieved for n if and only if equalities hold both in (1) and in $e_3 \leq e(G)$. This implies that $e(G) = 12(n - 2)/5$ for n if and only if G is a connected Θ_4 -free plane graph on n vertices such that each edge in G belongs to one 3-face and one 4-face. We next construct such an extremal graph for n and Θ_4 . Let $n = 20k + 12$ for some integer $k \geq 0$. Let G_0 be the graph depicted in Figure 1(a), we then construct G_k on n vertices recursively for all $k \geq 1$ via the illustration given in Figure 1(b): the entire graph G_{k-1} is placed into the center quadrangle of Figure 1(b), and the entire G_0 is then placed between the two given bold quadrangles of Figure 1(b) (in such a way that these are identified with the bold quadrangles of Figure 1(a)). One can check that G_k is Θ_4 -free with $n = 20k + 12$ vertices and $12(n - 2)/5$ edges for all $k \geq 0$. ■

We next prove that $ex_{\mathcal{P}}(n, \Theta_5) \leq 5(n-2)/2$ and then demonstrate that this bound is tight, in the sense that there are infinitely many values of n for which it is attained exactly.

Theorem 2.3 $ex_{\mathcal{P}}(n, \Theta_5) \leq 5(n-2)/2$ for all $n \geq 5$, with equality when $n \equiv 50 \pmod{120}$.

Proof. Let G be a Θ_5 -free plane graph on $n \geq 5$ vertices. We show by induction on n that $e(G) \leq 5(n-2)/2$. The statement is trivially true when $n = 5$ because any Θ_5 -free plane graph on five vertices has at most seven edges. So we may assume that $n \geq 6$. Next assume that there exists a vertex $u \in V(G)$ with $d_G(u) \leq 2$. By the induction hypothesis, $e(G \setminus u) \leq 5(n-3)/2$ and so $e(G) = e(G \setminus u) + d_G(u) \leq 5(n-3)/2 + 2 < 5(n-2)/2$, as desired. So we may assume that $\delta(G) \geq 3$. Assume next that G is disconnected. Let $G_1, \dots, G_s, G_{s+1}, \dots, G_{s+t}$ be all components of G such that $|G_1| = \dots = |G_s| = 4$ and $5 \leq |G_{s+1}| \leq \dots \leq |G_{s+t}|$, where $s \geq 0$ and $t \geq 0$ are integers with $s+t \geq 2$ and $4s + |G_{s+1}| + \dots + |G_{s+t}| = n$. Then $e(G_i) = 6$ for all $i \in [s]$ because $\delta(G) \geq 3$, and $e(G_j) \leq 5(|G_j| - 2)/2$ for all $j \in \{s+1, \dots, s+t\}$ by the induction hypothesis. Therefore,

$$\begin{aligned} e(G) &\leq 6s + \frac{5(|G_{s+1}| + \dots + |G_{s+t}| - 2t)}{2} \\ &= \frac{5(n-2)}{2} - \frac{(8(s+t) + 2t - 10)}{2} < \frac{5(n-2)}{2}, \end{aligned}$$

as desired. So we may further assume that G is connected.

Since G is a connected plane graph on $n \geq 6$ vertices, we see that G has no face of size at most two. Hence

$$2e(G) = 3f_3 + 4f_4 + \sum_{i \geq 5} if_i \geq 3f_3 + 4f_4 + 5(f - f_3 - f_4) = 5f - 2f_3 - f_4, \quad (2)$$

which implies that $f \leq (2e(G) + 2f_3 + f_4)/5$. Note that no 3-face in G has its three edges in $E_{3,3}$ because G is Θ_5 -free and $n \geq 6$. It follows that $e_{3,3} \leq f_3$. By Observation 2.1(b),

$$3f_3 = e_3 + e_{3,3} \leq e_3 + f_3 \text{ and so } f_3 \leq e_3/2. \quad (3)$$

It is worth noting that a 4-face and a 3-face in G cannot have exactly one edge in common, else G would contain Θ_5 as a subgraph. Since $\delta(G) \geq 3$, we see that a 4-face and a 3-face in G cannot have exactly two edges in common. Hence, every 4-face and every 3-face in G have no edge in common and so $E_{3,4} = \emptyset$. Thus, $e_3 + e_4 \leq e(G)$. By Observation 2.1(a,b), $e_{4,4} \leq e_4$ and $4f_4 = e_4 + e_{4,4}$. It follows that

$$4f_4 \leq 2e_4 \leq 2(e(G) - e_3) \text{ and so } f_4 \leq (e(G) - e_3)/2. \quad (4)$$




Figure 2: Construction of G_k .

Now with the last inequalities in (3) and (4), and the fact that $f \leq (2e(G) + 2f_3 + f_4)/5$ and $e_3 \leq e(G)$, we obtain $f \leq 3e(G)/5$. By Euler's formula, $n - 2 = e(G) - f \geq 2e(G)/5$. Hence $e(G) \leq 5(n - 2)/2$, as desired.

From the proof above, we see that equality in $e(G) \leq 5(n - 2)/2$ is achieved for n if and only if equalities hold in (2), (3) and (4) and in $e_3 \leq e(G)$. This implies that $e(G) = 5(n - 2)/2$ for n if and only if G is a connected Θ_5 -free plane graph on n vertices satisfying: each 3-face in G has exactly two edges in $E_{3,3}$; each edge in G belongs to either one 3-face and one 5-face or two 3-faces. We next construct such an extremal plane graph for n and Θ_5 . Let $n = 120k + 50$ for some integer $k \geq 0$. Let G_0 be the graph depicted in Figure 2(a), we then construct G_k of order n recursively for all $k \geq 1$ via the illustration given in Figure 2(b): the entire graph G_{k-1} is placed into the center pentagon of Figure 2(b), and the entire G_0 is then placed between the two given bold pentagons of Figure 2(b) (in such a way that these are identified with the bold pentagons of Figure 2(a)). One can check that G_k is Θ_5 -free with $n = 120k + 50$ vertices and $5(n - 2)/2$ edges for all $k \geq 0$. ■

Finally, we prove an upper bound for $ex_p(n, \Theta_6)$ in Theorem 2.4. Figure 3 illustrates all possible graphs for which equality in Theorem 2.4 is attained when $n = 9$. However, we shall see in Corollary 2.5 that equality is not possible for all $n \geq 10$.

Theorem 2.4 $ex_p(n, \Theta_6) \leq 18(n - 2)/7$ for all $n \geq 6$, with equality when $n = 9$.

Proof. Let G be an extremal plane graph for Θ_6 and $n \geq 6$. We shall prove that $e(G) \leq 18(n - 2)/7$ by induction on n . When $n = 6$, we show that $e(G) \leq 10$. Suppose that $e(G) \geq 11$.




Figure 3: All possible graphs achieving equality in Theorem 2.4 and Corollary 2.5 when $n = 9$.

Then G is isomorphic to either a plane triangulation on six vertices or a plane triangulation on six vertices with one edge removed. Note that all plane triangulations on 6 vertices are depicted in Figure 4. It is easy to check that G has a Hamiltonian cycle and so G contains a graph in Θ_6 as subgraph, a contradiction. Hence, $e(G) \leq 10 < 18(n-2)/7$ when $n = 6$. So we may assume that $n \geq 7$. Next assume that there exists a vertex $u \in V(G)$ with $d_G(u) \leq 2$. By the induction hypothesis, $e(G \setminus u) \leq 18(n-3)/7$ and so $e(G) = e(G \setminus u) + d_G(u) \leq 18(n-3)/7 + 2 < 18(n-2)/7$, as desired. So we may assume that $\delta(G) \geq 3$. Assume next that G is disconnected. Then each component of G has exactly four, five or at least six vertices because $\delta(G) \geq 3$. Let $G_1, \dots, G_r, G_{r+1}, \dots, G_{r+s}, G_{r+s+1}, \dots, G_{r+s+t}$ be all components of G such that

$$|G_1| = \dots = |G_r| = 4, |G_{r+1}| = \dots = |G_{r+s}| = 5, \text{ and } 6 \leq |G_{r+s+1}| \leq \dots \leq |G_{r+s+t}|,$$

where $r, s, t \geq 0$ are integers with $r + s + t \geq 2$ and $4r + 5s + |G_{r+s+1}| + \dots + |G_{r+s+t}| = n$. Since G is an extremal plane graph for Θ_6 , we see that $e(G_i) = 6$ for all $i \in [r]$ and $e(G_j) = 9$ for all $j \in \{r+1, \dots, r+s\}$. By the induction hypothesis, $e(G_k) \leq 18(|G_k| - 2)/7$ for all $k \in \{r+s+1, \dots, r+s+t\}$. Therefore,

$$\begin{aligned} e(G) &\leq 6r + 9s + \frac{18(|G_{r+s+1}| + \dots + |G_{r+s+t}| - 2t)}{7} \\ &= \frac{18(n-2)}{7} - \frac{(27(r+s+t) + 3r + 9t - 36)}{7} \\ &< \frac{18(n-2)}{7}, \end{aligned}$$

as desired. So we may assume that G is connected.




Figure 4: All plane triangulations on 6 vertices.

Next assume that G contains a cut-vertex, say u . Let H be a smallest component of $G \setminus u$, and let $G_1 := G[V(H) \cup \{u\}]$ and $G_2 := G \setminus V(H)$. Then $|G_1| \leq |G_2|$ and $|G_1| + |G_2| = n + 1$. Since $\delta(G) \geq 3$, we see that $4 \leq |G_1| \leq |G_2|$. Assume first that $|G_2| \leq 5$. Then $e(G_i) \leq 3|G_i| - 6$ for all $i \in \{1, 2\}$. Hence, $e(G) = e(G_1) + e(G_2) \leq 3(|G_1| + |G_2|) - 12 = 3n - 9 \leq 18(n - 2)/7$ because $n \leq 9$, with equality when both G_1 and G_2 are isomorphic to K_5 minus one edge, and so G is isomorphic to the graphs depicted in Figure 3. Assume next that $|G_2| \geq 6$. Then $e(G_2) \leq 18(|G_2| - 2)/7$ by the induction hypothesis. Note that $e(G_1) \leq 3|G_1| - 6$ when $|G_1| \leq 5$ and $e(G_1) \leq 18(|G_1| - 2)/7$ when $|G_1| \geq 6$ by the induction hypothesis. Therefore, when $|G_1| \leq 5$,

$$\begin{aligned} e(G) &= e(G_1) + e(G_2) \leq 3|G_1| - 6 + \frac{18(n + 1 - |G_1| - 2)}{7} \\ &= \frac{18(n - 2)}{7} - \frac{(24 - 3|G_1|)}{7} < \frac{18(n - 2)}{7}; \end{aligned}$$

when $|G_1| \geq 6$,

$$e(G) = e(G_1) + e(G_2) \leq \frac{18(|G_1| + |G_2| - 4)}{7} < \frac{18(n - 2)}{7}.$$

So we may assume that G is 2-connected. By Observation 2.1(d) and the fact that $\delta(G) \geq 3$, each face in G is bounded by a cycle.




Figure 5: Three possible configurations of H with $v_2v_3 \in E_{4,4} \cup E_{3,5}$.

Assume next that $E_{4,4} \cup E_{3,5} \neq \emptyset$. Let $v_2v_3 \in E_{4,4} \cup E_{3,5}$. Since $\delta(G) \geq 3$, let F_1 and F_2 be the two faces of G having v_2v_3 in common such that the size of F_1 is at least the size of F_2 . Then G must contain a plane subgraph H isomorphic to the graphs depicted in Figure 5(a,b) when $v_2v_3 \in E_{4,4}$, and in Figure 5(c) when $v_2v_3 \in E_{3,5}$, because G is Θ_6 -free. Let H_1 and H_2 be the induced plane subgraphs of G with boundary v_1, v_2, v_5 and v_1, v_3, v_4 , respectively. Then $|H_1| + |H_2| = n + 1$, and $|H_i| \geq 6$ for all $i \in [2]$ because G is Θ_6 -free, 2-connected and $\delta(G) \geq 3$. By the induction hypothesis, $e(H_i) \leq 18(|H_i| - 2)/7$ for all $i \in [2]$. Thus,

$$e(G) = e(H_1) + e(H_2) + |\{v_2v_3\}| < 18(n - 2)/7.$$

We may now further assume that $E_{4,4} \cup E_{3,5} = \emptyset$. Then $e_{4,4} = 0$ and $e_{3,5} = 0$.

It is easy to see that G is not a plane triangulation and so $\sum_{i \geq 4} f_i \geq 0$. We next show that $\sum_{i \geq 5} f_i \neq 0$. Suppose $\sum_{i \geq 5} f_i = 0$. Then $f_3 + f_4 = f$ and $f_4 > 0$. Note that $e_{4,4} = 0$. It follows that every edge of a 4-face of G belongs to $E_{3,4}$, and so G contains a Θ_6 subgraph, a contradiction. Thus $\sum_{i \geq 5} f_i \neq 0$. We may further assume that the outer face of G is neither a 3-face nor a 4-face. Then

$$\begin{aligned} 2e(G) &= 3f_3 + 4f_4 + 5f_5 + \sum_{i \geq 6} if_i \\ &\geq 3f_3 + 4f_4 + 5f_5 + 6(f - f_3 - f_4 - f_5) \\ &= 6f - 3f_3 - 2f_4 - f_5, \end{aligned} \tag{5}$$

which implies that $6f \leq 2e(G) + 3f_3 + 2f_4 + f_5$.




Figure 6: All possible configurations of H_F , where all dashed edges are in $E_{3,3}$, and no solid edges are in $E_{3,3}$.

We next find an upper bound for each of f_3, f_4 and f_5 . To get an upper bound for f_3 , we first show that $5e_{3,3} \leq 6f_3$. Let F be a 3-face in G with $|E(F) \cap E_{3,3}| \geq 1$. Clearly, $|E(F) \cap E_{3,3}| \leq 3$. Since G is Θ_6 -free and the outer face of G is not a 3-face, there exists a plane subgraph H_F of G with $|H_F| \leq 5$ such that F is a face (not the outer face) of H_F ; all faces of H_F , except the outer face of H_F and any face of H_F that is not a face in G , are 3-faces; and no edges on the boundary of the outer face of H_F and any face of H_F that is not a face in G are in $E_{3,3}$. The possible configurations of H_F are shown in Figure 6. When H_F is isomorphic to the graph depicted in Figure 6(b), H_F contains six edges in $E_{3,3}$ and five 3-faces of G . From all possible configurations of H_F , we see that $e_{3,3} \leq 6f_3/5$. Hence,

$$3f_3 = e_3 + e_{3,3} \leq e_3 + 6f_3/5, \text{ and so } f_3 \leq 5e_3/9. \tag{6}$$

To get an upper bound for f_4 , we next show that $4f_4 \leq 2(e(G) - e_3)$. By Observation 2.1(b,c),

$$4f_4 = e_4 \text{ and } e(G) \geq e_3 + e_4 - e_{3,4}. \tag{7}$$

We next show that $e_{3,4} \leq e(G) - e_3$.

This is trivially true when $e_{3,4} = 0$. Assume that $e_{3,4} \neq 0$. Let F and F' be a 4-face and a 3-face in G , respectively, such that F and F' share an edge in common. We may assume that F

has vertices v_1, v_2, v_3, v_4 in order and F' has vertices v_1, v_4, v_5 in order. Note that F and F' are not outer face in G . Observe that if $v_i v_{i+1}$ belongs to $E_{3,4}$ for any $i \in \{1, 2, 3\}$, then $v_i v_{i+1}$ belongs to the 4-face F and the 3-face with vertices v_i, v_{i+1}, v_5 in order, else G would not be Θ_6 -free. Since $n \geq 7$, there exists some $k \in \{1, 2, 3\}$ such that $v_k v_{k+1} \notin E_{3,4}$. Then $v_k v_{k+1} \in E_{4,j}$ for some $j \geq 5$ because $e_{4,4} = 0$. We next show that F has at most two edges in $E_{3,4}$. Suppose $|E(F) \cap E_{3,4}| = 3$. We may assume that $k = 2$. Then $v_1 v_2, v_3 v_4 \in E_{3,4}$. Thus $v_1 v_2$ belongs to the 4-face F and the 3-face with vertices v_1, v_2, v_5 in order; and $v_3 v_4$ belongs to the 4-face F and the 3-face with vertices v_3, v_4, v_5 in order. Since G is Θ_6 -free, we see that $v_5 v_2 \in E_{3,j}$ for some $j \geq 6$, $v_5 v_3 \in E_{3,j}$ for some $j \geq 6$, and $v_2 v_3 \in E_{4,j}$ for some $j \geq 6$. But then $G + v_2 v_4$ is Θ_6 -free, contrary to the choice of G . Thus F has at most two edges in $E_{3,4}$, and so F has at least two edges not in E_3 . This holds for each 4-face in G . Hence, $e_{3,4} \leq e(G) - e_3$.

By (7),

$$4f_4 = e_4 \leq e(G) - e_3 + e_{3,4} \leq 2(e(G) - e_3). \quad (8)$$

Note that $e_{3,5} = 0$. By Observation 2.1(a), $e_{5,5} \leq e_5 \leq e(G) - e_3$. By Observation 2.1(b),

$$5f_5 = e_5 + e_{5,5} \leq 2(e(G) - e_3). \quad (9)$$

Combining $e_3 \leq e(G)$ with the upper bounds on f_3, f_4, f_5 given in (6), (8), (9), we have

$$\begin{aligned} 6f &\leq 2e(G) + 3f_3 + 2f_4 + f_5 \\ &\leq 2e(G) + 5e_3/3 + (e(G) - e_3) + 2(e(G) - e_3)/5 \\ &= 17e(G)/5 + 4e_3/15 \\ &\leq 11e(G)/3. \end{aligned}$$

It follows that $f \leq 11e(G)/18$. By Euler's formula, $n - 2 \geq e(G) - f \geq 7e(G)/18$. Hence $e(G) \leq 18(n - 2)/7$, as desired. \blacksquare

Corollary 2.5 *Let K_5^- be the graph obtained from K_5 by deleting one edge. Then*

(a) $ex_{\mathcal{P}}(n, \Theta_6 \cup \{K_5^-\}) \leq 12(n - 2)/5$ for all $n \geq 7$.

(b) $ex_{\mathcal{P}}(n, \Theta_6) < 18(n - 2)/7$ for all $n \geq 10$.

Proof. To prove (a), let G be an extremal plane graph for $n \geq 7$ and $\Theta_6 \cup \{K_5^-\}$. We prove that $e(G) \leq 12(n - 2)/5$ by induction on n . Since G is Θ_6 -free, by Theorem 2.4, $e(G) \leq \lfloor 18(n - 2)/7 \rfloor = 12(n - 2)/5$ when $n = 7$. So we may assume that $n \geq 8$. Similar to the proof of Theorem 2.4, we see that $e_{3,3} \leq f_3$, because G is K_5^- -free and so no H_F is isomorphic to the graphs depicted in

Figure 6(b, c). By Observation 2.1(b), $f_3 \leq e_3/2$. This, together with the upper bounds for f_4, f_5 given in (8) and (9), implies that

$$\begin{aligned}
6f &\leq 2e(G) + 3f_3 + 2f_4 + f_5 \\
&\leq 2e(G) + 3e_3/2 + e(G) - e_3 + 2(e(G) - e_3)/5 \\
&= 17e(G)/5 + e_3/10 \\
&\leq 7e(G)/2.
\end{aligned}$$

It follows that $f \leq 7e(G)/12$. By Euler's formula, $n - 2 \geq e(G) - f \geq 5e(G)/12$. Hence $e(G) \leq 12(n - 2)/5$.




Figure 7: An example of constructing the graph G' from a graph G .

To prove (b), let G be an extremal plane graph for Θ_6 and $n \geq 10$. By Theorem 2.4, $e(G) \leq 18(n - 2)/7$. Suppose $e(G) = 18(n - 2)/7$. By Corollary 2.5(a), G is not K_5^- -free. From the proof of Theorem 2.4, we see that equality in $e(G) \leq 18(n - 2)/7$ is achieved for n if and only if all the equalities hold in (5), (6), (8), (9) and in $e_3 \leq e(G)$. This implies that $e(G) = 18(n - 2)/7$ for n if and only if G is a 2-connected Θ_6 -free plane graph on n vertices satisfying: G consists entirely of K_5^- 's and 6-faces, no two K_5^- 's share an edge, and no two 6-faces have an edge in common. Let G' be the graph obtained from G by deleting the two vertices not on the outer face in each $H_F = K_5^-$, an example is shown in Figure 7. Then G' consists of 3-faces and 6-faces such that each edge of G' belongs to one 3-face and one 6-face. Clearly, G' is 2-connected because G is 2-connected. Let f'_i be the number of i -faces in G' . Let $f' = \sum_{i \geq 1} f'_i$. Then $3f'_3 = e(G') = 6f'_6$ and $f' = f'_3 + f'_6$. Thus $|G'| - 2 = e(G') - f' = e(G')/2$ and so $e(G') = 2|G'| - 4$, which implies that $\delta(G') \leq 3$. Since G' is 2-connected, we have $\delta(G') \geq 2$. Note that each vertex of G' must have even degree because the adjacent faces are alternatively of size 3 and size 6. Thus $\delta(G') = 2$. Let $v \in V(G')$ be a vertex of degree two in G' . Let u_1vu_2 and $u_1vu_2u_3u_4u_5$ be the vertices in order on the boundary of the two adjacent faces containing v , respectively. Then $G'[\{u_1, v, u_2, u_3, u_4, u_5\}]$ contains a graph in Θ_6 as a subgraph. Thus G is not Θ_6 -free, a contradiction. ■

It is worth noting that every C_6 -free graph is certainly Θ_6 -free. Hence, $ex_{\mathcal{P}}(n, C_6) \leq ex_{\mathcal{P}}(n, \Theta_6)$. Corollary 2.6 follows immediately from Theorem 2.4.

Corollary 2.6 $ex_{\mathcal{P}}(n, C_6) \leq 18(n - 2)/7$ for all $n \geq 6$, with equality when $n = 9$.

Acknowledgments.

The authors would like to thank the referees for their careful reading and helpful comments.

Zi-Xia Song would like to thank Yongtang Shi and the Chern Institute of Mathematics at Nankai University for hospitality and support during her visit in May 2017.

This work was partially supported by National Natural Science Foundation of China, Natural Science Foundation of Tianjin (No. 17JCQNJC00300), the China-Slovenia bilateral project “Some topics in modern graph theory” (No. 12-6), Open Project Foundation of Intelligent Information Processing Key Laboratory of Shanxi Province (No. CICIP2018005), and the Fundamental Research Funds for the Central Universities, Nankai University.

References

- [1] B. Bollobás, *Modern Graph Theory*, Springer, 2013.
- [2] G. A. Dirac, Some theorems on abstract graphs, *Proc. London Math. Soc.* 2 (1952) 69–81.
- [3] C. Dowden, Extremal C_4 -free/ C_5 -free planar graphs, *J. Graph Theory* 83 (2016) 213–230.
- [4] P. Erdős and A. H. Stone, On the structure of linear graphs, *Bull. Amer. Math. Soc.* 52 (1946) 1087–1091.
- [5] Z. Füredi, Turán type problems, “*Surveys in Combinatorics*”, London Math. Soc. Lecture Note Ser. 166, Cambridge Univ. Press, Cambridge, 1991, pp. 253–300.
- [6] Z. Füredi and T. Jiang, Hypergraph Turán numbers of linear cycles, *J. Combin. Theory Ser. A* 123 (2014) 252–270.
- [7] Z. Füredi, T. Jiang and R. Seiver, Exact solution of the hypergraph Turán problem for k -uniform linear paths, *Combinatorica* 34 (2014) 299–322.
- [8] M. Horňák, S. Jendrol’, I. Schiermeyer and R. Soták, Rainbow numbers for cycles in plane triangulations, *J. Graph Theory* 78 (2015) 248–257.
- [9] P. Keevash, Hypergraph Turán problems, “*Surveys in Combinatorics 2011*”, London Math. Soc. Lecture Note Ser. 392, Cambridge Univ. Press, Cambridge (2011), pp. 83–139.
- [10] A. Kostochka, D. Mubayi and J. Verstraëte, Turán problems and shadows I: Paths and cycles, *J. Combin. Theory Ser. A* 129 (2015) 57–79.
- [11] Y. Lan, Y. Shi and Z-X. Song, Planar anti-Ramsey numbers of paths and cycles, to appear in *Discrete Mathematics*.
- [12] P. Turán, Eine Extremalaufgabe aus der Graphentheorie, (*Hungarian*) *Mat. Fiz. Lapok* 48 (1941) 436–452.