

The Real-rootedness of Generalized Narayana Polynomials related to the Boros-Moll Polynomials

Herman Z.Q. Chen¹, Arthur L.B. Yang², Philip B. Zhang³

^{1,2}Center for Combinatorics, LPMC
Nankai University, Tianjin 300071, P. R. China

³College of Mathematical Science
Tianjin Normal University, Tianjin 300387, P. R. China

Email: ¹zqchern@163.com, ²yang@nankai.edu.cn, ³zhangbiaonk@163.com

Abstract. In this paper, we prove the real-rootedness of a family of generalized Narayana polynomials, which arose in the study of the infinite log-concavity of the Boros-Moll polynomials. We establish certain recurrence relations for these Narayana polynomials, from which we derive the real-rootedness. To prove the real-rootedness, we use a sufficient condition, due to Liu and Wang, to determine whether two polynomials have interlaced zeros. The recurrence relations are verified with the help of the Mathematica package *HolonomicFunctions*.

AMS Classification 2010: 05A15, 26C10

Keywords: Generalized Narayana polynomials, real zeros, interlacing zeros.

1 Introduction

For any nonnegative integers n and m , let

$$N_n(x) = \frac{1}{n+1} \sum_{k=0}^n \binom{n+1}{k} \binom{n+1}{k+1} x^k, \quad (1)$$

$$N_{n,m}(x) = \sum_{k=0}^n \left(\binom{n}{k} \binom{m}{k} - \binom{n}{k+1} \binom{m}{k-1} \right) x^k. \quad (2)$$

The polynomial $N_n(x)$ is the classical Narayana polynomial. It is well known that $N_n(x)$ has only real zeros. Moreover, it is easy to verify that both $N_{n,n}(x)$ and $N_{n+1,n}(x)$ are just the polynomial $N_n(x)$. While, it seems that the polynomials $N_{n,m}(x)$ were not well studied for general n and m . In this paper, we shall prove that the polynomials $N_{n,m}(x)$ have only real zeros for any n and m . Let us first review some backgrounds of the polynomials $N_{n,m}(x)$.

The polynomials $N_{n,m}(x)$ arose in the study of the infinite log-concavity of the Boros-Moll polynomials. The Boros-Moll polynomials were introduced by Boros and Moll [1]

in their study of a quartic integral, and they obtained the following expression for the Boros-Moll polynomials:

$$P_n(x) = 2^{-2n} \sum_j 2^j \binom{2n-2j}{n-j} \binom{n+j}{j} (x+1)^j.$$

Recall that a finite nonnegative sequence $\{a_k\}_{k=0}^n$ is said to be log-concave if

$$a_k^2 - a_{k+1}a_{k-1} \geq 0, \quad \text{for } 0 \leq k \leq n,$$

where, for convenience, we set $a_{-1} = 0$ and $a_{n+1} = 0$. We say that it is infinitely log-concave if for any $i \geq 1$ the i -th iterative sequence $\{\mathcal{L}^i(a_k)\}_{k=0}^n$ is nonnegative, where \mathcal{L} is the operator acting on $\{a_k\}_{k=0}^n$ as follows

$$\mathcal{L}(a_k) = a_k^2 - a_{k+1}a_{k-1}, \quad \text{for } 0 \leq k \leq n.$$

We say that a polynomial

$$f(x) = \sum_{k=0}^n a_k x^k$$

is infinitely log-concave if its coefficient sequence $\{a_k\}_{k=0}^n$ is infinitely log-concave. Boros and Moll proposed the following conjecture.

Conjecture 1.1 ([1]). *The polynomial $P_n(x)$ is infinitely log-concave.*

The log-concavity of $P_n(x)$ was first conjectured by Moll [11], and then was proved by Kauers and Paule [7]. The 2-fold log-concavity of $P_n(x)$ was proved by Chen and Xia [5]. Brändén [2] proposed an innovative approach to the higher-fold log-concavity of $P_n(x)$. He conjectured the real-rootedness of some variations of $P_n(x)$, from which its 3-fold log-concavity can be deduced. Brändén's conjectures were later confirmed by Chen, Dou and Yang [4]. While Conjecture 1.1 is open, Brändén [2] has proved the infinite log-concavity of real-rooted polynomials, which was independently conjectured by Stanley, McNamara and Sagan [10], and Fisk [6].

Theorem 1.2 ([2]). *If*

$$f(x) = \sum_{k=0}^n a_k x^k$$

is a real-rooted polynomial with nonnegative coefficients, then so is the polynomial

$$\sum_{k=0}^n (a_k^2 - a_{k-1}a_{k+1})x^k.$$

The well known Newton's inequality states that if a polynomial $f(x)$ has only real zeros, then it must be log-concave. Therefore, Theorem 1.2 implies the infinite log-concavity of the real-rooted polynomials. Motivated by Brändén's theorem, we are led to study the real-rootedness of the following polynomial:

$$Q_n(x) = \sum_{k=0}^n (d_k(n)^2 - d_{k-1}(n)d_{k+1}(n))x^k,$$

where

$$d_k(n) = 2^{-2n} \sum_{j=k}^n 2^j \binom{2n-2j}{n-j} \binom{n+j}{j} \binom{j}{k}$$

is the coefficient of x^k in the Boros-Moll polynomial $P_n(x)$. We have the following conjecture.

Conjecture 1.3. *For any $n \geq 1$, the polynomial $Q_n(x)$ has only real zeros.*

Since the log-concavity of $P_n(x)$ is known, by Theorem 1.2, Conjecture 1.3 would imply Conjecture 1.1. Note that the polynomial $Q_n(x)$ may be rewritten as

$$Q_n(x) = \sum_{i=0}^n \sum_{j=0}^n 2^{i+j} \binom{2n-2i}{n-i} \binom{2n-2j}{n-j} \binom{n+i}{i} \binom{n+j}{j} N_{i,j}(x),$$

where $N_{i,j}(x)$ is the Narayana polynomial defined by (2). The numerical evidence suggests that the polynomial $N_{n,m}(x)$ has only real zeros for any n and m . Our main result is as follows.

Theorem 1.4. *For any $m, n \geq 0$, the polynomial $N_{m,n}(x)$ has only real zeros.*

The remainder of this paper is organized as follows. In the next section, we shall give an overview of some tools which will be used to prove Theorem 1.4. In Section 3, we shall establish some interlacing property concerning the polynomials $N_{n,m}(x)$, from which we derive Theorem 1.4.

2 Preliminaries

The results contained in this section serve as a reference point used in Section 3.

Let us first introduce the definition of interlacing. Given two real-rooted polynomials $f(x)$ and $g(x)$ with positive leading coefficients, We say that $g(x)$ *interlaces* $f(x)$, denoted $g(x) \preceq f(x)$, if

$$\cdots \leq s_2 \leq r_2 \leq s_1 \leq r_1,$$

where $\{r_i\}$ and $\{s_j\}$ are the sets of zeros of $f(x)$ and $g(x)$, respectively.

Liu and Wang [9] obtained the following sufficient condition to determine whether two polynomials have interlaced zeros.

Theorem 2.1 ([9, Theorem 2.3]). *Let $F(x), f(x), g_1(x), \dots, g_k(x)$ be polynomials with real coefficients satisfying the following conditions.*

(a) *There exist some polynomials $\phi(x), \varphi_1(x), \dots, \varphi_k(x)$ with real coefficients such that*

$$F(x) = \phi(x)f(x) + \varphi_1(x)g_1(x) + \dots + \varphi_k(x)g_k(x) \quad (3)$$

and $\deg F(x) = \deg f(x)$ or $\deg F(x) = \deg f(x) + 1$;

(b) *The polynomials $f(x), g_1(x), \dots, g_k(x)$ are real-rooted polynomials, and moreover $g_j(x) \preceq f(x)$ for each $1 \leq j \leq k$;*

(c) *The leading coefficients of $F(x)$ and $g_j(x)$ have the same sign for each $1 \leq j \leq k$.*

Suppose that $\varphi_j(r) \leq 0$ for each j and each zero r of $f(x)$. Then $F(x)$ has only real zeros and $f(x) \preceq F(x)$.

We shall use the above result to prove the real-rootedness of $N_{n,m}(x)$. The key point is to prove certain recurrence relations related to these polynomials. As will be shown later, the coefficients of these recurrence relations look complicated. Thanks to Zeilberger's holonomic systems approach to special function identities, Koutschan (private communication) pointed out that these recurrence relations can be easily verified by using the Mathematica package *HolonomicFunctions*, see [8, 13]. The Ore algebras introduced in [12] serve as a unifying framework to represent such recurrence relations. These algebras were obtained by applying Ore extensions to some base rings, also called Ore polynomial rings. Let S_n denote the shift operator with respect to n . Let $\mathbb{R}(n, x)$ denote the field of rational functions in n and x over the field \mathbb{R} of real numbers. The Ore algebra used throughout this paper could be considered as $\mathbb{R}(n, x)\langle S_n \rangle$, which consists of all linear operators of the form $\sum_{i=0}^r p_i S_n^i$, where $r \geq 0$ and $p_i \in \mathbb{R}(n, x)$. Suppose that the polynomial sequence $\{f_n(x)\}_{n \geq 0}$ satisfies certain recurrence relation $\sum_{i=0}^k a_i f_{i+n}(x) = 0$, where $a_i \in \mathbb{R}(n, x)$, and then the Ore polynomial of such a recurrence relation is given by $\sum_{i=0}^k a_i S_n^i$. For each $f \in \mathbb{R}(n, x)$, the annihilator of f with respect to $\mathbb{R}(n, x)\langle S_n \rangle$ is defined by

$$\text{Ann}_{\mathbb{R}(n, x)\langle S_n \rangle}(f) = \{P \in \mathbb{R}(n, x)\langle S_n \rangle \mid P(f) = 0\},$$

which is a left ideal in $\mathbb{R}(n, x)\langle S_n \rangle$. For more information on the Ore algebras and the Ore polynomials, see Koutschan [8] and Ore [12].

To be self-contained, we give an example to illustrate the use of this package. It is well known that the classical Narayana polynomials $N_n(x)$ given in (1) satisfy the following recurrence relation:

$$(n+3)N_{n+1}(x) = (2n+3)(x+1)N_n(x) - n(x-1)^2N_{n-1}(x).$$

This can be proved in the following way by using the package:

1. Convert the above recurrence to an Ore polynomial in the Ore algebra:

$$\text{In[1]:= rec = ToOrePolynomial}[(2 * n + 3) * (x + 1) * f[n] - n * (x - 1)^2 * f[n - 1] - (n + 3) * f[n + 1], f[n]]$$

$$\text{Out[1]= } \{(4 + n)S_n^2 + (-5 - 2n - 5x - 2nx)S_n + (1 + n - 2x - 2nx + x^2 + nx^2)\}$$

2. Generate a (Gröebner) basis of the annihilator A of the input (i.e., the set of all recurrence/differential relations that the input satisfies) using the command `Annihilator`:

$$\text{In[2]:= ann = Annihilator[Sum[Binomial[n + 1, k] * Binomial[n + 1, k + 1] * x^k / (n + 1), {k, 0, n}], S[n]]$$

$$\text{Out[2]= } \{(4 + n)S_n^2 + (-5 - 2n - 5x - 2nx)S_n + (1 + n - 2x - 2nx + x^2 + nx^2)\}$$

3. Reduce the Ore polynomial rec modulo A using the command `OreReduce`. If it returns 0, then rec is an element of the set A and hence the recurrence relation is valid.

$$\text{In[3]:= OreReduce[rec, ann]$$

$$\text{Out[3]= } 0$$

3 Proof

The objective of this section is to give a proof of Theorem 1.4, namely the real-rootedness of the polynomial $N_{n,m}(x)$. We first derive certain recurrence relations for these polynomials. For nonnegative integers t and n , let

$$\underline{N}_n^{(t)}(x) = N_{n,n+t}(x), \quad \overline{N}_n^{(t)}(x) = N_{n+t,n}(x). \quad (4)$$

The polynomials $\underline{N}_n^{(t)}(x)$ satisfy the following recurrence relation.

Theorem 3.1. *For nonnegative integers t and $n \geq 1$, we have*

$$\underline{N}_{n+1}^{(t)}(x) = \frac{a_0 + a_1x + a_2x^2}{(n + t + 3)(n + 1)(c_0 + c_1x)} \underline{N}_n^{(t)}(x) - \frac{n(n + t)(x - 1)^2(b_0 + b_1x)}{(n + t + 3)(n + 1)(c_0 + c_1x)} \underline{N}_{n-1}^{(t)}(x), \quad (5)$$

where

$$\begin{aligned}
a_0 &= -(2n^3 + (2t + 5)n^2 + (2t + 3)n), \\
a_1 &= (2t(t + 2)n^3 + 3t(t + 2)^2n^2 + (t(t + 2)(t^2 + 5t + 5))n \\
&\quad + (t(t + 1)(t + 2)(t + 3)/2)), \\
a_2 &= (t + 1)((2t + 2)n^3 + (3t^2 + 9t + 5)n^2 + (2t + 3)(t^2 + 3t + 1)n \\
&\quad + t(t + 1)(t + 2)(t + 3)/2), \\
b_0 &= -(n + 1), \\
b_1 &= (t + 1)^2n + (t + 1)(t^2 + 4t + 2)/2, \\
c_0 &= -n, \\
c_1 &= (t + 1)^2n + t(t + 1)(t + 2)/2.
\end{aligned}$$

Proof. We shall prove an equivalent form of this recurrence relation, which is obtained by multiplying $(n + t + 3)(n + 1)(c_0 + c_1x)$ on both sides of (5). This could be converted into an Ore polynomial as follows:

$$\begin{aligned}
\text{In[4]:= } \mathbf{rec} &= \mathbf{ToOrePolynomial}[(\mathbf{a0} + \mathbf{a1} * \mathbf{x} + \mathbf{a2} * \mathbf{x}^2) * \mathbf{f}[n] - (\mathbf{n} * (\mathbf{n} + \mathbf{t}) * (\mathbf{x} - \\
&\quad \mathbf{1})^2 * (\mathbf{b0} + \mathbf{b1} * \mathbf{x})) * \mathbf{f}[n - 1] - (\mathbf{n} + \mathbf{t} + \mathbf{3}) * (\mathbf{n} + \mathbf{1}) * (\mathbf{c0} + \mathbf{c1} * \mathbf{x}) * \mathbf{f}[n + \\
&\quad \mathbf{1}] /. \mathbf{MapThread}[\mathbf{Rule}, \{\{\mathbf{a0}, \mathbf{a1}, \mathbf{a2}, \mathbf{b0}, \mathbf{b1}, \mathbf{c0}, \mathbf{c1}\}, \{-(2 * \mathbf{n}^3 + (2 * \mathbf{t} + 5) * \mathbf{n}^2 + \\
&\quad (2 * \mathbf{t} + 3) * \mathbf{n}), 2 * \mathbf{t} * (\mathbf{t} + 2) * \mathbf{n}^3 + 3 * \mathbf{t} * (\mathbf{t} + 2)^2 * \mathbf{n}^2 + (\mathbf{t} * (\mathbf{t} + 2) * (\mathbf{t}^2 + 5 * \mathbf{t} + 5)) * \\
&\quad \mathbf{n} + \mathbf{t} * (\mathbf{t} + 1) * (\mathbf{t} + 2) * ((\mathbf{t} + 3) / 2), (\mathbf{t} + 1) * ((2 * \mathbf{t} + 2) * \mathbf{n}^3 + (3 * \mathbf{t}^2 + 9 * \mathbf{t} + 5) * \mathbf{n}^2 + \\
&\quad (2 * \mathbf{t} + 3) * (\mathbf{t}^2 + 3 * \mathbf{t} + 1) * \mathbf{n} + \mathbf{t} * (\mathbf{t} + 1) * (\mathbf{t} + 2) * ((\mathbf{t} + 3) / 2)), -(\mathbf{n} + \mathbf{1}), (\mathbf{t} + \mathbf{1})^2 * \\
&\quad \mathbf{n} + (\mathbf{t} + \mathbf{1}) * ((\mathbf{t}^2 + 4 * \mathbf{t} + 2) / 2), -\mathbf{n}, (\mathbf{t} + \mathbf{1})^2 * \mathbf{n} + \mathbf{t} * (\mathbf{t} + \mathbf{1}) * ((\mathbf{t} + 2) / 2)\}], \mathbf{f}[n]];
\end{aligned}$$

Then compute a (Gröebner) basis ann of the set of all recurrence/differential relations that $\underline{N}_n^{(t)}(x)$ satisfies, and reduce the Ore polynomial rec modulo ann :

$$\begin{aligned}
\text{In[5]:= } \mathbf{ann} &= \mathbf{Annihilator}[\mathbf{Sum}[(\mathbf{Binomial}[n, k] * \mathbf{Binomial}[n + t, k] - \mathbf{Binomial}[n, k + \\
&\quad \mathbf{1}] * \mathbf{Binomial}[n + t, k - 1]) * \mathbf{x}^k, \{k, 0, n\}], \mathbf{S}[n]]; \\
\text{In[6]:= } &\mathbf{OreReduce}[\mathbf{rec}, \mathbf{ann}] \\
\text{Out[6]= } &0
\end{aligned}$$

We have the desired output. This completes the proof. \square

Next we come to proving the real-rootedness of $\underline{N}_n^{(t)}(x)$.

Theorem 3.2. *For any $t \geq 0$ and $n \geq 0$, the polynomial $\underline{N}_n^{(t)}(x)$ has only real zeros, and moreover, we have $\underline{N}_n^{(t)}(x) \preceq \underline{N}_{n+1}^{(t)}(x)$.*

Proof. We use induction on n . It is straightforward to verify that

$$\underline{N}_0^{(t)}(x) = 1, \quad \underline{N}_1^{(t)}(x) = 1 + (t + 1)x, \quad \underline{N}_0^{(t)}(x) \preceq \underline{N}_1^{(t)}(x).$$

Assume that $\underline{N}_{n-1}^{(t)}(x) \preceq \underline{N}_n^{(t)}(x)$. We see that the recurrence relation (5) is of the form (3) in Theorem 2.1 with $k = 1$, where

$$\begin{aligned} F(x) &= \underline{N}_{n+1}^{(t)}(x), \\ f(x) &= \underline{N}_n^{(t)}(x), \\ g_1(x) &= \underline{N}_{n-1}^{(t)}(x), \\ \phi(x) &= \frac{a_0 + a_1x + a_2x^2}{(n+t+3)(n+1)(c_0 + c_1x)}, \\ \varphi_1(x) &= -\frac{n(n+t)(x-1)^2(b_0 + b_1x)}{(n+t+3)(n+1)(c_0 + c_1x)}. \end{aligned}$$

Here, $a_0, a_1, a_2, b_0, b_1, c_0, c_1$ are given by (5). Note that for any $n, t \geq 0$ the coefficients of $\underline{N}_{n+1}^{(t)}(x)$ are nonnegative, since, for any $0 \leq k \leq n$, the coefficient of x^k in $\underline{N}_{n+1}^{(t)}(x)$ is

$$\begin{aligned} [x^k]\underline{N}_{n+1}^{(t)}(x) &= \binom{n}{k} \binom{n+t}{k} - \binom{n}{k+1} \binom{n+t}{k-1} \\ &= \left(1 - \frac{k}{k+1} \cdot \frac{n-k}{n-k+t+1}\right) \binom{n}{k} \binom{n+t}{k} > 0. \end{aligned}$$

It is clear that for any $x \leq 0$, we have $\varphi_1(x) \leq 0$. By Theorem 2.1, the polynomial $\underline{N}_{n+1}^{(t)}(x)$ is real-rooted, and moreover $\underline{N}_n^{(t)}(x) \preceq \underline{N}_{n+1}^{(t)}(x)$. \square

We have the following recurrence relation for $\overline{N}_n^{(t)}(x)$.

Theorem 3.3. *For nonnegative integers t and $n \geq 1$, we have*

$$\overline{N}_{n+1}^{(t)}(x) = \frac{a_0 + a_1x + a_2x^2}{(n+t+1)(n+3)(c_0 + c_1x)} \overline{N}_n^{(t)}(x) - \frac{n(n+t)(x-1)^2(b_0 + b_1x)}{(n+t+1)(n+3)(c_0 + c_1x)} \overline{N}_{n-1}^{(t)}(x), \quad (6)$$

where

$$\begin{aligned} a_0 &= -(2n+3)(n+t)(n+t+1), \\ a_1 &= 3t(t-2)(t+1)^2/2 + t(t-2)(t^2+7t+5)n \\ &\quad + 3t(t-2)(t+2)n^2 + 2t(t-2)n^3, \\ a_2 &= t^2(t-1)(t+1)^2/2 + (t-1)(2t^3+3t^2+t-3)n \\ &\quad + (t-1)(3t^2+3t-5)n^2 + 2(t-1)^2n^3, \\ b_0 &= -n-1-t, \\ b_1 &= (t-1)^2n + (t-1)t^2/2 + (t-1)^2, \\ c_0 &= -n-t, \\ c_1 &= (t-1)^2n + (t-1)t^2/2. \end{aligned}$$

Proof. The proof is similar to that of Lemma 3.1. We need to prove an equivalent form of (6) obtained by multiplying $(n+t+1)(n+3)(c_0+c_1x)$ on both sides. This could be converted into an Ore polynomial as follows:

```

In[7]:= rec = ToOrePolynomial[(a0 + a1 * x + a2 * x^2) * f[n] - (n * (n + t) * (x -
1)^2 * (b0 + b1 * x)) * f[n - 1] - (n + 3) * (n + t + 1) * (c0 + c1 * x) * f[n +
1] /. MapThread[Rule, {{a0, a1, a2, b0, b1, c0, c1}, {-(2 * n + 3) * (n + t) * (n +
t + 1), 3 * t * (t - 2) * (t + 1)^2/2 + t * (t - 2) * (t^2 + 7t + 5) * n + 3 * t * (t - 2) *
(t + 2) * n^2 + 2 * t * (t - 2) * n^3, t^2 * (t - 1) * (t + 1)^2/2 + (t - 1) * (2 * t^3 + 3 * t^2 +
t - 3) * n + (t - 1) * (3 * t^2 + 3 * t - 5) * n^2 + 2 * (t - 1)^2 * n^3, -n - 1 - t, (t -
1)^2 * n + (t - 1) * t^2/2 + (t - 1)^2, -n - t, (t - 1)^2 * n + (t - 1) * t^2/2}}], f[n]];

```

Then compute a (Gröebner) basis ann of the set of all recurrence/differential relations that $\overline{N}_n^{(t)}(x)$ satisfies, and reduce the Ore polynomial rec modulo ann :

```

In[8]:= ann = Annihilator[Sum[(Binomial[n + t, k] * Binomial[n, k] - Binomial[n + t, k +
1] * Binomial[n, k - 1]) * x^k, {k, 0, n + t}], S[n]];
In[9]:= OreReduce[rec, ann]
Out[9]= 0

```

The output is 0, as desired. This completes the proof. \square

We now prove the real-rootedness of $\overline{N}_n^{(t)}(x)$.

Theorem 3.4. *For any $n, t \geq 0$, the polynomial $\overline{N}_n^{(t)}(x)$ has only real zeros. If $t \geq 2$, then $\overline{N}_n^{(t)}(x)$ has one and only one positive zero.*

Proof. Note that both the polynomials $\overline{N}_n^{(0)}(x)$ and $\overline{N}_n^{(1)}(x)$ are the classical Narayana polynomial, which is known to be real-rooted.

We proceed to consider the case of $t \geq 2$. We first prove that $\overline{N}_n^{(t)}(x)$ has one and only one positive zero. Note that for any $n \geq 0$ and $t \geq 2$, $\overline{N}_n^{(t)}(x)$ is polynomial in x of degree $n+1$, and for any $0 \leq k \leq n+1$, the coefficient of x^k in $\overline{N}_n^{(t)}(x)$ is

$$\binom{n+t}{k} \binom{n}{k} - \binom{n+t}{k+1} \binom{n}{k-1} = \frac{n+1-kt}{(n+1)(k+1)} \binom{n+t}{k} \binom{n+1}{k}.$$

Therefore, the number of changes in sign of the coefficients is 1. By Descartes' rule, the polynomial $\overline{N}_n^{(t)}(x)$ has at most one positive zero. Moreover, we see that

$$[x^0]\overline{N}_n^{(t)}(x) = 1 > 0, \quad [x^{n+1}]\overline{N}_n^{(t)}(x) = -\binom{n+t}{n+2} < 0.$$

Thus, the polynomial $\overline{N}_n^{(t)}(x)$ has one and only one positive zero.

Next we claim that $\overline{N}_n^{(t)}(x)$ has n negative zeros, and moreover, for any $n \geq 1$,

$$r_{n+1}^{(n+1)} < r_n^{(n)} < r_n^{(n+1)} < r_{n-1}^{(n)} < \cdots < r_2^{(n)} < r_2^{(n+1)} < r_1^{(n)} < r_1^{(n+1)} < 0,$$

where $\{r_k^{(n)}\}_{k=0}^n$ and $\{r_k^{(n+1)}\}_{k=0}^{n+1}$ are the negative zeros of $\overline{N}_n^{(t)}(x)$ and $\overline{N}_{n+1}^{(t)}(x)$ respectively.

Before proving the above claim, let us note the following property: for any $x < 0$, $n \geq 1$ and $t \geq 2$, clearly we have

$$-\frac{n(n+t)(x-1)^2(b_0+b_1x)}{(n+t+1)(n+3)(c_0+c_1x)} < 0.$$

To prove the claim, we use induction on n . Let us first prove the base case of $n = 1$. We already showed that $\overline{N}_1^{(t)}(x)$ has one and only one positive zero. Since $\overline{N}_1^{(t)}(x)$ is of degree 2 and $[x^0]\overline{N}_1^{(t)}(x) = 1$, it also has one negative zero $r_1^{(1)}$. By the recurrence (6), we see that $\overline{N}_2^{(t)}(r_1^{(1)}) < 0$ since $\overline{N}_0^{(t)}(r_1^{(1)}) > 0$. Moreover, we have $\overline{N}_2^{(t)}(0) = 1 > 0$ and $\overline{N}_2^{(t)}(-\infty) > 0$. Thus, $\overline{N}_2^{(t)}(x)$ has two negative zeros $r_1^{(2)}, r_2^{(2)}$ and moreover $r_2^{(2)} < r_1^{(1)} < r_1^{(2)} < 0$, as claimed.

Assume that the claim is true for n . We proceed to show that it is also true for $n + 1$. From (6) we deduce that

$$(-1)^k \overline{N}_{n+1}^{(t)}(r_k^{(n)}) > 0, \quad \text{for any } 1 \leq k \leq n.$$

Moreover, we have $\overline{N}_{n+1}^{(t)}(0) = 1 > 0$ and $(-1)^{n+1} \overline{N}_{n+1}^{(t)}(-\infty) > 0$. Thus, the polynomial $\overline{N}_{n+1}^{(t)}(x)$ has $n + 1$ negative zeros $\{r_k^{(n+1)}\}_{k=0}^{n+1}$, and moreover, for each $1 \leq k \leq n$, we have $r_{k+1}^{(n+1)} < r_k^{(n)} < r_k^{(n+1)}$, as claimed. This completes the proof. \square

Combining Theorems 3.2 and 3.4, we complete the proof of Theorem 1.4.

Acknowledgements. We would like to thank the referee for the valuable comments which helped to improve the manuscript. We also would like to thank Dr. Christoph Koutschan for helpful suggestions on the use of the Mathematica package *Holonomic-Functions*.

The first and second authors were supported by the National Science Foundation of China. The third author was partially supported by NSFC grant 11626172 and PHD Program 52XB1616 of Tianjin Normal University.

References

- [1] G. Boros and V. Moll, *Irresistible integrals*, Cambridge Univ. Press, Cambridge, 2004.

- [2] P. Brändén, Iterated sequences and the geometry of zeros, *J. Reine Angew. Math.* **658** (2011), 115–131.
- [3] F. Brenti, Unimodal, log-concave and Pólya frequency sequences in combinatorics, *Mem. Amer. Math. Soc.* **81** (1989), no. 413, viii+106 pp.
- [4] W. Y. C. Chen, D. Q. J. Dou and A. L. B. Yang, Brändén’s conjectures on the Boros-Moll polynomials, *Int. Math. Res. Not.* **20** (2013), 4819–4828.
- [5] W. Y. C. Chen and E. X. W. Xia, 2-log-concavity of the Boros-Moll polynomials, *Proc. Edinb. Math. Soc.* (2) **56** (2013), no. 3, 701–722.
- [6] S. Fisk, Questions about determinants and polynomials, arXiv:0808.1850.
- [7] M. Kauers and P. Paule, A computer proof of Moll’s log-concavity conjecture, *Proc. Amer. Math. Soc.* **135** (2007), no. 12, 3847–3856.
- [8] C. Koutschan, *Advanced Applications of the Holonomic Systems Approach*, PhD thesis, RISC, Johannes Kepler University, Linz, Austria, 2009.
- [9] L. L. Liu and Y. Wang, A unified approach to polynomial sequences with only real zeros, *Adv. Appl. Math.* **38** (2007), no. 4, 542–560.
- [10] P. R. W. McNamara and B. E. Sagan, Infinite log-concavity: developments and conjectures, *Adv. Appl. Math.* **44** (2010), no. 1, 1–15.
- [11] V. H. Moll, The evaluation of integrals: a personal story, *Notices Amer. Math. Soc.* **49** (2002), no. 3, 311–317.
- [12] O. Ore, Theory of non-commutative polynomials, *Annals Math.* **34** (1933), no. 3, 480–508.
- [13] D. Zeilberger, A holonomic systems approach to special functions identities, *J. Comput. Appl. Math.* **32** (1990), no. 3, 321–368.