

3 GRAPHS WITH LARGE GENERALIZED (EDGE-)CONNECTIVITY

4 XUELIANG LI

5 *Center for Combinatorics and LPMC*
6 *Nankai University*
7 *Tianjin 300071, China*

8 **e-mail:** lxl@nankai.edu.cn

9 AND

10 YAPING MAO

11 *Department of Mathematics*
12 *Qinghai Normal University*
13 *Qinghai 810008, China*

14 **e-mail:** maoyaping@ymail.com

15 **Abstract**

16 The generalized k -connectivity $\kappa_k(G)$ of a graph G , introduced by Hager
17 in 1985, is a nice generalization of the classical connectivity. Recently,
18 as a natural counterpart, we proposed the concept of generalized k -edge-
19 connectivity $\lambda_k(G)$. In this paper, graphs of order n such that $\kappa_k(G) =$
20 $n - \frac{k}{2} - 1$ and $\lambda_k(G) = n - \frac{k}{2} - 1$ for even k are characterized.

21 **Keywords:** (edge-)connectivity; Steiner tree; internally disjoint trees; edge-
22 disjoint trees; packing; generalized (edge-)connectivity..

23 **2010 Mathematics Subject Classification:** Type 2010 Mathematics
24 05C40, 05C05, 05C70, 05C75.

25 1. INTRODUCTION

26 All graphs considered in this paper are undirected, finite and simple. We refer
27 to the book [3] for graph theoretical notation and terminology not described here.
28 For a graph G , let $V(G)$, $E(G)$, \overline{G} denote the set of vertices, the set of edges of
29 G and the complement, respectively. Let $d_G(v)$ denote the degree of the vertex
30 v in G . As usual, the *union* of two graphs G and H is the graph, denoted by

31 $G \cup H$, with vertex set $V(G) \cup V(H)$ and edge set $E(G) \cup E(H)$. Let mH be the
 32 disjoint union of m copies of a graph H . If M is a subset of edges of a graph G ,
 33 the subgraph of G induced by M is denoted by $G[M]$, and $G - M$ denotes the
 34 subgraph obtained by deleting the edges of M from G . If $M = \{e\}$, we simply
 35 write $G - e$ for $G - \{e\}$. If $S \subseteq V(G)$, the subgraph of G induced by S is denoted
 36 by $G[S]$. For $S \subseteq V(G)$, we denote $G - S$ the subgraph obtained by deleting the
 37 vertices of S together with the edges incident with them from G . We denote by
 38 $E_G[X, Y]$ the set of edges of G with one end in X and the other end in Y . If
 39 $X = \{x\}$, we simply write $E_G[x, Y]$ for $E_G[\{x\}, Y]$. A subset M of $E(G)$ is called
 40 a *matching* of G if the edges of M satisfy that no two of them are adjacent in G .
 41 A matching M saturates a vertex v , or v is said to be *M -saturated*, if some edge
 42 of M is incident with v ; otherwise, v is *M -unsaturated*. If every vertex of G is
 43 M -saturated, the matching M is *perfect*. M is a *maximum matching* if G has no
 44 matching M' with $|M'| > |M|$.

45 Connectivity and edge-connectivity are two of the most basic concepts of
 46 graph-theoretic subjects, both in a combinatorial sense and an algorithmic sense.
 47 As we know, the classical connectivity has two equivalent definitions. The *con-*
 48 *nectivity* of a graph G , written $\kappa(G)$, is the minimum size of a set $S \subseteq V(G)$ such
 49 that $G - S$ is disconnected or has only one vertex. If $G - S$ is disconnected we
 50 call such a set S a *vertex cut-set* for G . We call this definition the ‘cut’ version
 51 definition of connectivity. A well-known Menger’s theorem provides an equiva-
 52 lent definition of connectivity, which can be called the ‘path’ version definition
 53 of connectivity. For any two distinct vertices x and y in G , the *local connectivity*
 54 $\kappa_G(x, y)$ is the maximum number of internally disjoint paths connecting x and
 55 y . Then $\kappa(G) = \min\{\kappa_G(x, y) \mid x, y \in V(G), x \neq y\}$ is defined to be the *con-*
 56 *nectivity* of G . Similarly, the classical edge-connectivity also has two equivalent
 57 definitions. The *edge-connectivity* of G , written $\lambda(G)$, is the minimum size of an
 58 edge set $M \subseteq E(G)$ such that $G - M$ is disconnected or has only one vertex.
 59 We call this definition the ‘cut’ version definition of edge-connectivity. Menger’s
 60 theorem also provides an equivalent definition of edge-connectivity, which can
 61 be called the ‘path’ version definition. For any two distinct vertices x and y in
 62 G , the *local edge-connectivity* $\lambda_G(x, y)$ is the maximum number of edge-disjoint
 63 paths connecting x and y . Then $\lambda(G) = \min\{\lambda_G(x, y) \mid x, y \in V(G), x \neq y\}$ is
 64 defined to be the *edge-connectivity* of G . For connectivity and edge-connectivity,
 65 Oellermann gave a survey paper on this subject, see [34].

66 Although there are many elegant and powerful results on connectivity in
 67 graph theory, the classical connectivity and edge-connectivity also have their
 68 defects. So people want some generalizations of both connectivity and edge-
 69 connectivity. For the ‘cut’ version definition of connectivity, we are looking for
 70 a minimum vertex-cut with no consideration about the number of components
 71 of $G - S$. Two graphs with the same connectivity may have different degrees of

72 vulnerability in the sense that the deletion of a vertex cut-set of minimum cardi-
 73 nality from one graph may produce a graph with considerably more components
 74 than in the case of the other graph. For example, the star $K_{1,n}$ and the path
 75 P_{n+1} ($n \geq 3$) are both trees of order $n + 1$ and therefore connectivity 1, but the
 76 deletion of a cut-vertex from $K_{1,n}$ produces a graph with n components while
 77 the deletion of a cut-vertex from P_{n+1} produces only two components. Char-
 78 trand et al. [4] generalized the ‘cut’ version definition of connectivity. For an
 79 integer k ($k \geq 2$) and a graph G of order n ($n \geq k$), the k -connectivity $\kappa'_k(G)$
 80 is the smallest number of vertices whose removal from G produces a graph with
 81 at least k components or a graph with fewer than k vertices. Thus, for $k = 2$,
 82 $\kappa'_2(G) = \kappa(G)$. For more details about k -connectivity, we refer to [4, 6, 35, 36].
 83 The k -edge-connectivity, which is a generalization of the ‘cut’ version definition
 84 of classical edge-connectivity was initially introduced by Boesch and Chen [2] and
 85 subsequently studied by Goldsmith in [7, 8] and Goldsmith et al. [9]. For more
 86 details, we refer to [1, 34].

87 The generalized connectivity of a graph G , introduced by Hager [12], is a
 88 natural and nice generalization of the ‘path’ version definition of connectivity.
 89 For a graph $G = (V, E)$ and a set $S \subseteq V$ of at least two vertices, an S -Steiner tree
 90 or a Steiner tree connecting S (or simply, an S -tree) is a subgraph $T = (V', E')$ of
 91 G that is a tree with $S \subseteq V'$. Two Steiner trees T and T' connecting S are said to
 92 be *internally disjoint* if $E(T) \cap E(T') = \emptyset$ and $V(T) \cap V(T') = S$. For $S \subseteq V(G)$
 93 and $|S| \geq 2$, the *generalized local connectivity* $\kappa(S)$ is the maximum number
 94 of internally disjoint Steiner trees connecting S in G . Note that when $|S| = 2$
 95 a minimal Steiner tree connecting S is just a path connecting the two vertices
 96 of S . For an integer k with $2 \leq k \leq n$, *generalized k -connectivity* (or *k -tree-*
 97 *connectivity*) is defined as $\kappa_k(G) = \min\{\kappa(S) \mid S \subseteq V(G), |S| = k\}$. Clearly, when
 98 $|S| = 2$, $\kappa_2(G)$ is nothing new but the connectivity $\kappa(G)$ of G , that is, $\kappa_2(G) =$
 99 $\kappa(G)$, which is the reason why one addresses $\kappa_k(G)$ as the generalized connectivity
 100 of G . By convention, for a connected graph G with less than k vertices, we set
 101 $\kappa_k(G) = 1$. Set $\kappa_k(G) = 0$ when G is disconnected. This concept appears to
 102 have been introduced by Hager in [12]. It is also studied in [5] for example,
 103 where the exact value of the generalized k -connectivity of complete graphs are
 104 obtained. Note that the generalized k -connectivity and the k -connectivity of a
 105 graph are indeed different. Take for example, the graph H_1 obtained from a
 106 triangle with vertex set $\{v_1, v_2, v_3\}$ by adding three new vertices u_1, u_2, u_3 and
 107 joining v_i to u_i by an edge for $1 \leq i \leq 3$. Then $\kappa_3(H_1) = 1$ but $\kappa'_3(H_1) = 2$.
 108 There are many results on the generalized connectivity or tree-connectivity, we
 109 refer to [5, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 37]. Apart from the concept
 110 of tree-connectivity, Hager also introduced another tree-connectivity parameter,
 111 called the *pendant tree-connectivity* of a graph in [12]. For the tree-connectivity,
 112 we only search for edge-disjoint trees which include S and are vertex-disjoint with

113 the exception of the vertices in S . But pendant tree-connectivity further requires
 114 the degree of each vertex of S in a Steiner tree connecting S equal to one. Note
 115 that it is a special case of the tree-connectivity.

116 As a natural counterpart of the generalized connectivity, we introduced in
 117 [32] the concept of generalized edge-connectivity, which is a generalization of the
 118 ‘path’ version definition of edge-connectivity. For $S \subseteq V(G)$ and $|S| \geq 2$, the
 119 *generalized local edge-connectivity* $\lambda(S)$ is the maximum number of edge-disjoint
 120 Steiner trees connecting S in G . For an integer k with $2 \leq k \leq n$, the *general-*
 121 *ized k -edge-connectivity* $\lambda_k(G)$ of G is then defined as $\lambda_k(G) = \min\{\lambda(S) \mid S \subseteq$
 122 $V(G) \text{ and } |S| = k\}$. It is also clear that when $|S| = 2$, $\lambda_2(G)$ is nothing new but
 123 the standard edge-connectivity $\lambda(G)$ of G , that is, $\lambda_2(G) = \lambda(G)$, which is the
 124 reason why we address $\lambda_k(G)$ as the generalized edge-connectivity of G . Also set
 125 $\lambda_k(G) = 0$ when G is disconnected. Results on the generalized edge-connectivity
 126 can be found in [28, 29, 32].

127 In fact, Mader [19] was studying an extension of Menger’s theorem to inde-
 128 pendent sets of three or more vertices. We know from Menger’s theorem that if
 129 $S = \{u, v\}$ is a set of two independent vertices in a graph G , then the maximum
 130 number of internally disjoint u - v paths in G equals the minimum number of ver-
 131 tices that separate u and v . For a set $S = \{u_1, u_2, \dots, u_k\}$ of k vertices ($k \geq 2$)
 132 in a graph G , an S -*path* is defined as a path between a pair of vertices of S that
 133 contains no other vertices of S . Two S -paths P_1 and P_2 are said to be *internally*
 134 *disjoint* if they are vertex-disjoint except for their endvertices. If S is a set of
 135 independent vertices of a graph G , then a vertex set $U \subseteq V(G)$ with $U \cap S = \emptyset$ is
 136 said to *totally separate* S if every two vertices of S belong to different components
 137 of $G - U$. Let S be a set of at least three independent vertices in a graph G .
 138 Let $\mu(G)$ denote the maximum number of internally disjoint S -paths and $\mu'(G)$
 139 the minimum number of vertices that totally separate S . A natural extension of
 140 Menger’s theorem may well be suggested, namely: If S is a set of independent
 141 vertices of a graph G and $|S| \geq 3$, then $\mu(S) = \mu'(S)$. However, the statement is
 142 not true in general. Take the above graph H_1 for example. For $S = \{v_1, v_2, v_3\}$,
 143 $\mu(S) = 1$ but $\mu'(S) = 2$. Mader proved that $\mu(S) \geq \frac{1}{2}\mu'(S)$. Moreover, the
 144 bound is sharp. Lovász conjectured an edge analogue of this result and Mader
 145 proved this conjecture and established its sharpness. For more details, we refer
 146 to [19, 20, 34].

147 In addition to being natural combinatorial measures, the Steiner Tree Pack-
 148 ing Problem and the generalized edge-connectivity can be motivated by their
 149 interesting interpretation in practice as well as theoretical consideration. From a
 150 theoretical perspective, both extremes of this problem are fundamental theorems
 151 in combinatorics. One extreme of the problem is when we have two terminal-
 152 s. In this case internally (edge-)disjoint trees are just internally (edge-)disjoint
 153 paths between the two terminals, and so the problem becomes the well-known

154 Menger theorem. The other extreme is when all the vertices are terminals. In
 155 this case internally disjoint Steiner trees and edge-disjoint trees are just edge-
 156 disjoint spanning trees of the graph, and so the problem becomes the classical
 157 Nash-Williams-Tutte theorem.

Theorem 1.1. (Nash-Williams [33], Tutte [39]) *A multigraph G contains a system of ℓ edge-disjoint spanning trees if and only if*

$$\|G/\mathcal{P}\| \geq \ell(|\mathcal{P}| - 1)$$

158 *holds for every partition \mathcal{P} of $V(G)$, where $\|G/\mathcal{P}\|$ denotes the number of cross-*
 159 *ing edges in G , i.e., edges between distinct parts of \mathcal{P} .*

160 The generalized edge-connectivity is related to an important problem, which
 161 is called the *Steiner Tree Packing Problem*. For a given graph G and $S \subseteq V(G)$,
 162 this problem asks to find a set of maximum number of edge-disjoint Steiner
 163 trees connecting S in G . One can see that the Steiner Tree Packing Problem
 164 studies local properties of graphs, but the generalized edge-connectivity focuses
 165 on global properties of graphs. The generalized edge-connectivity and the Steiner
 166 Tree Packing Problem have applications in *VLSI* circuit design, see [10, 11, 38].
 167 In this application, a Steiner tree is needed to share an electronic signal by a
 168 set of terminal nodes. Another application, which is our primary focus, arises
 169 in the Internet Domain. Imagine that a given graph G represents a network.
 170 We choose arbitrary k vertices as nodes. Suppose that one of the nodes in G
 171 is a *broadcaster*, and all the other nodes are either *users* or *routers* (also called
 172 *switches*). The broadcaster wants to broadcast as many streams of movies as
 173 possible, so that the users have the maximum number of choices. Each stream of
 174 movie is broadcasted via a tree connecting all the users and the broadcaster. So,
 175 in essence we need to find the maximum number of Steiner trees connecting all
 176 the users and the broadcaster, namely, we want to get $\lambda(S)$, where S is the set
 177 of the k nodes. Clearly, it is a Steiner tree packing problem. Furthermore, if we
 178 want to know whether for any k nodes the network G has the above properties,
 179 then we need to compute $\lambda_k(G) = \min\{\lambda(S)\}$ in order to prescribe the reliability
 180 and the security of the network.

181 The following two observations are easily seen from the definitions.

182 **Observation 1.2.** *Let k, n be two integers with $3 \leq k \leq n$. For a connected*
 183 *graph G of order n , $\kappa_k(G) \leq \lambda_k(G) \leq \delta(G)$.*

184 **Observation 1.3.** *Let k, n be two integers with $3 \leq k \leq n$. If H is a spanning*
 185 *subgraph of G of order n , then $\lambda_k(H) \leq \lambda_k(G)$.*

186 Chartrand et al. in [5] got the exact value of the generalized k -connectivity
 187 for the complete graph K_n .

188 **Lemma 1.4.** [5] *For every two integers n and k with $2 \leq k \leq n$, $\kappa_k(K_n) =$
189 $n - \lceil k/2 \rceil$.*

190 In [32] we obtained some results on the generalized k -edge-connectivity. The
191 following results are restated, which will be used later.

192 **Lemma 1.5.** [32] *For every two integers n and k with $2 \leq k \leq n$, $\lambda_k(K_n) =$
193 $n - \lceil k/2 \rceil$.*

194 **Lemma 1.6.** [32] *Let k, n be two integers with $3 \leq k \leq n$. For a connected graph
195 G of order n , $1 \leq \kappa_k(G) \leq \lambda_k(G) \leq n - \lceil k/2 \rceil$. Moreover, the upper and lower
196 bounds are sharp.*

197 We also characterized graphs attaining the upper bound and obtained the
198 following result.

199 **Lemma 1.7.** [32] *Let k, n be two integers with $3 \leq k \leq n$. For a connected graph
200 G of order n , $\kappa_k(G) = n - \lceil \frac{k}{2} \rceil$ or $\lambda_k(G) = n - \lceil \frac{k}{2} \rceil$ if and only if $G = K_n$ for even
201 k ; $G = K_n - M$ for odd k , where M is a set of edges such that $0 \leq |M| \leq \frac{k-1}{2}$.*

202 One may notice that the graphs with $\kappa_k(G) = n - \lceil \frac{k}{2} \rceil$ are the same as the
203 graphs with $\lambda_k(G) = n - \lceil \frac{k}{2} \rceil$. Our motivation of this paper is to ask whether
204 the graphs with $\kappa_k(G) = n - \lceil \frac{k}{2} \rceil - 1$ are different from the graphs with $\lambda_k(G) =$
205 $n - \lceil \frac{k}{2} \rceil - 1$. In this paper, graphs of order n such that $\kappa_k(G) = n - \lceil \frac{k}{2} \rceil - 1$ and
206 $\lambda_k(G) = n - \lceil \frac{k}{2} \rceil - 1$ for any even k are characterized.

207 **Theorem 1.8.** *Let n and k be two integers such that k is even and $4 \leq k \leq n$,
208 and G be a connected graph of order n . Then $\kappa_k(G) = n - \frac{k}{2} - 1$ if and only
209 if $G = K_n - M$ where M is a set of edges such that $1 \leq \Delta(K_n[M]) \leq \frac{k}{2}$ and
210 $1 \leq |M| \leq k - 1$.*

211 The above result can also be established for the generalized k -edge-connectivity,
212 which is stated as follows.

213 **Theorem 1.9.** *Let n and k be two integers such that k is even and $4 \leq k \leq n$,
214 and G be a connected graph of order n . Then $\lambda_k(G) = n - \frac{k}{2} - 1$ if and only if
215 $G = K_n - M$ where M is a set of edges satisfying one of the following conditions:*

- 216 (1) $\Delta(K_n[M]) = 1$ and $1 \leq |M| \leq \lfloor \frac{n}{2} \rfloor$;
217 (2) $2 \leq \Delta(K_n[M]) \leq \frac{k}{2}$ and $1 \leq |M| \leq k - 1$.

218 2. MAIN RESULT

219 To begin with, we give the following lemmas.

220 **Lemma 2.1.** *If G is a graph obtained from the complete graph K_n by deleting a*
 221 *set of edges M such that $\Delta(K_n[M]) \geq r$, then $\lambda_k(G) \leq n - 1 - r$.*

222 **Proof.** Since $\Delta(K_n[M]) \geq r$, there exists at least one vertex, say v , such that
 223 $d_{K_n[M]}(v) \geq r$. Then $d_G(v) = n - 1 - d_{K_n[M]}(v) \leq n - 1 - r$. So $\delta(G) \leq d_G(v) \leq$
 224 $n - 1 - r$. From Observation 1.2, $\lambda_k(G) \leq \delta(G) \leq n - 1 - r$. ■

225 **Corollary 2.2.** *For every two integers n and k with $4 \leq k \leq n$, if k is even and*
 226 *M is a set of edges in the complete graph K_n such that $\Delta(K_n[M]) \geq \frac{k}{2} + 1$, then*
 227 *$\kappa_k(K_n - M) \leq \lambda_k(K_n - M) < n - \frac{k}{2} - 1$.*

228 **Remark 1.** From Corollary 2.2, if $\kappa_k(K_n - M) = n - \frac{k}{2} - 1$ or $\lambda_k(K_n - M) =$
 229 $n - \frac{k}{2} - 1$ for k even, then $\Delta(K_n[M]) \leq \frac{k}{2}$.

230 In [32], we stated a useful lemma for general k .

231 Let $S \subseteq V(G)$ be such that $|S| = k$, and \mathcal{T} be a maximum set of edge-
 232 disjoint S -Steiner trees in G . Let \mathcal{T}_1 be the set of trees in \mathcal{T} whose edges belong
 233 to $E(G[S])$, and \mathcal{T}_2 be the set of S -Steiner trees containing at least one edge of
 234 $E_G[S, \bar{S}]$, where $\bar{S} = V(G) - S$. Thus, $\mathcal{T} = \mathcal{T}_1 \cup \mathcal{T}_2$ (Throughout this paper, \mathcal{T} ,
 235 \mathcal{T}_1 , \mathcal{T}_2 are defined in this way).

236 **Lemma 2.3.** [32] *Let G be a connected graph of order n , and $S \subseteq V(G)$ with*
 237 *$|S| = k$ ($3 \leq k \leq n$) and let T be a S -Steiner tree. If $T \in \mathcal{T}_1$, then T contains*
 238 *exactly $k - 1$ edges of $E(G[S])$. If $T \in \mathcal{T}_2$, then T contains at least k edges of*
 239 *$E(G[S]) \cup E_G[S, \bar{S}]$.*

240 **Lemma 2.4.** *For every two integers n and k with $4 \leq k \leq n$, if k is even and M*
 241 *is a set of edges of the complete graph K_n such that $|M| \geq k$ and $\Delta(K_n[M]) \geq 2$,*
 242 *then $\lambda_k(K_n - M) < n - \frac{k}{2} - 1$.*

243 **Proof.** Set $G = K_n - M$. We claim that there is an $S \subseteq V(G)$ with $|S| = k$ such
 244 that $|M \cap (E(K_n[S]) \cup E_{K_n}[S, \bar{S}])| \geq k$ and $|M \cap E(K_n[S])| \geq 1$. Choose a subset
 245 M' of M such that $|M'| = k$. Suppose that $K_n[M']$ contains s independent edges
 246 and r connected components C_1, \dots, C_r such that $\Delta(C_i) \geq 2$ ($1 \leq i \leq r$). Set
 247 $|V(C_i)| = n_i$ and $|E(C_i)| = m_i$. Then $m_i \geq n_i - 1$. For each C_i ($1 \leq i \leq r$), we
 248 select one of the vertices having maximum degree, say u_i . Set $X_i = V(C_i) - u_i$.

249 If there exists some X_j such that $|E(K_n[X_j])| \geq 1$, then we choose $X_i \subseteq S$
 250 for all $1 \leq i \leq r$. Since $|V(C_i)| = n_i$ and $X_i = V(C_i) - u_i$, we have $|X_i| = n_i - 1$.
 251 By such a choosing, the number of the vertices belonging to S is $\sum_{i=1}^r |X_i| =$
 252 $\sum_{i=1}^r (n_i - 1) \leq \sum_{i=1}^r m_i \leq k - s$. In addition, we select one endvertex of each
 253 independent edge into S . Till now, the total number of the vertices belonging to
 254 S is $\sum_{i=1}^r |X_i| + s \leq (k - s) + s = k$. Note that if $\sum_{i=1}^r |X_i| + s < k$, then we can
 255 add some other vertices in G into S such that $|S| = k$. Thus all edges of $E(C_i)$
 256 and the s independent edges are put into $E(K_n[S]) \cup E_{K_n}[S, \bar{S}]$, that is, all edges

257 of M' belong to $E(K_n[S]) \cup E_{K_n}[S, \bar{S}]$. So $|M \cap (E(K_n[S]) \cup E_{K_n}[S, \bar{S}])| \geq k$, as
 258 desired. Since $|E(K_n[X_j])| \geq 1$, it follows that $|M \cap (E(K_n[S]))| \geq 1$, as desired.

259 Suppose that $|E(K_n[X_i])| = 0$ for all $1 \leq i \leq r$. Then each C_i must be a
 260 star such that $|E(C_i)| \geq 2$. Recall that u_i is one of the vertices having maximum
 261 degree in C_i . Select one vertex from $V(C_i) - u_i$, say v_i . Put all the vertices of
 262 $Y_i = V(C_i) - v_i$ into S , that is, $Y_i \subseteq S$. Thus $|Y_i| = n_i - 1$. In addition, we
 263 choose one endvertex of each independent edge into S . By such a choosing, the
 264 total number of the vertices belonging to S is $\sum_{i=1}^r |Y_i| + s = \sum_{i=1}^r (n_i - 1) + s \leq$
 265 $\sum_{i=1}^r m_i + s \leq (k - s) + s = k$. Note that if $\sum_{i=1}^r |X_i| + s < k$ then we can add
 266 some other vertices in G into S such that $|S| = k$. Thus all edges of $E(C_i)$ and
 267 the s independent edges are put into $E(K_n[S]) \cup E_{K_n}[S, \bar{S}]$, that is, and all edges
 268 of M' belong to $E(K_n[S]) \cup E_{K_n}[S, \bar{S}]$. So $|M \cap (E(K_n[S]) \cup E_{K_n}[S, \bar{S}])| \geq k$,
 269 as desired. Since $|E(C_i)| \geq 2$, it follows that there is an edge $u_i w_i \in M \cap K_n[S]$
 270 where $w_i \in V(C_i) - \{u_i, v_i\}$, which implies that $|M \cap (E(K_n[S]))| \geq 1$, as desired.

271 From the above arguments, we conclude that there exists an $S \subseteq V(G)$ with
 272 $|S| = k$ such that $|M \cap (E(K_n[S]) \cup E_{K_n}[S, \bar{S}])| \geq k$ and $|M \cap (E(K_n[S]))| \geq 1$.
 273 Since each tree $T \in \mathcal{T}_1$ uses $k - 1$ edges in $E(G[S]) \cup E_G[S, \bar{S}]$, it follows that
 274 $|\mathcal{T}_1| \leq \binom{k}{2} - 1 / (k - 1) = \frac{k}{2} - \frac{1}{k-1}$, which results in $|\mathcal{T}_1| \leq \frac{k}{2} - 1$ since $|\mathcal{T}_1|$
 275 is an integer. From Lemma 2.3, each tree $T \in \mathcal{T}_2$ uses at least k edges of
 276 $E(G[S]) \cup E_G[S, \bar{S}]$. Thus $|\mathcal{T}_1|(k - 1) + |\mathcal{T}_2|k \leq |E(G[S])| + |E_G[S, \bar{S}]|$, that is,
 277 $|\mathcal{T}_1|k + |\mathcal{T}_2|k \leq |\mathcal{T}_1| + \binom{k}{2} + k(n - k) - k$. So $\lambda_k(G) = |\mathcal{T}| = |\mathcal{T}_1| + |\mathcal{T}_2| \leq$
 278 $n - \frac{k}{2} - 1 - \frac{1}{k} < n - \frac{k}{2} - 1$. \blacksquare

279 **Remark 2.** From Lemmas 1.7 and 2.4, if $\kappa_k(K_n - M) = n - \frac{k}{2} - 1$ or $\lambda_k(K_n - M) =$
 280 $n - \frac{k}{2} - 1$ for k even and $2 \leq \Delta(K_n[M]) \leq \frac{k}{2}$, then $1 \leq |M| \leq k - 1$, where
 281 $M \subseteq E(K_n)$.

282 **Lemma 2.5.** For every two integers n and k with $4 \leq k \leq n$, if k is even and M
 283 is a set of edges in the complete graph K_n such that $|M| \geq k$ and $\Delta(K_n[M]) = 1$,
 284 then $\kappa_k(K_n - M) < n - \frac{k}{2} - 1$.

285 **Proof.** Let $G = K_n - M$. Since $\Delta(K_n[M]) = 1$, it follows that M is a matching
 286 in K_n . Since $|M| \geq k$, we can choose $M_1 \subseteq M$ such that $|M_1| = k$. Let
 287 $M_1 = \{u_i w_i | 1 \leq i \leq k\}$. Choose $S = \{u_1, u_2, \dots, u_k\}$. We will show that
 288 $\kappa(S) < n - \frac{k}{2} - 1$. Clearly, $|\bar{S}| = n - k$, and let $\bar{S} = \{w_1, w_2, \dots, w_{n-k}\}$. Since
 289 each tree in \mathcal{T}_2 contains at least one vertex of \bar{S} , it follows that $|\mathcal{T}_2| \leq n - k$.
 290 By the definition of \mathcal{T}_1 , we have $|\mathcal{T}_1| \leq \frac{k}{2}$. If $|\mathcal{T}_1| \leq \frac{k}{2} - 2$, then $\kappa(S) \leq \lambda(S) =$
 291 $|\mathcal{T}| = |\mathcal{T}_1| + |\mathcal{T}_2| \leq (\frac{k}{2} - 2) + (n - k) = n - \frac{k}{2} - 2 < n - \frac{k}{2} - 1$, as desired. Let
 292 us assume $\frac{k}{2} - 1 \leq |\mathcal{T}_1| \leq \frac{k}{2}$.

293 Consider the case $|\mathcal{T}_1| = \frac{k}{2} - 1$. Recall that $|\mathcal{T}_2| \leq n - k$. Furthermore,
 294 we claim that $|\mathcal{T}_2| \leq n - k - 1$. Assume, to the contrary, that $|\mathcal{T}_2| = n - k$.
 295 Let T_1, T_2, \dots, T_{n-k} be the $n - k$ edge-disjoint S -Steiner trees in \mathcal{T}_2 . For each

296 tree T_i ($1 \leq i \leq n - k$), this tree only occupy one vertex of \bar{S} , say w_i . Since
 297 $u_i w_i \in M_1$ ($1 \leq i \leq k$), namely, $u_i w_i \notin E(G)$, and each T_i ($1 \leq i \leq k$) is an
 298 S -Steiner tree in \mathcal{T}_2 , it follows that this tree T_i must contain at least one edge
 299 in $G[S] = K_k$. So the trees T_1, T_2, \dots, T_k must use at least k edges in $G[S]$,
 300 and $|\mathcal{T}_1| = \frac{\binom{k}{2} - k}{k-1} = \frac{k-2}{2} - \frac{1}{k-1}$. Since $|\mathcal{T}_1|$ is an integer, we have $|\mathcal{T}_1| < \frac{k-2}{2}$,
 301 a contradiction. We conclude that $|\mathcal{T}_2| \leq n - k - 1$, and hence $\kappa(S) \leq \lambda(S) =$
 302 $|\mathcal{T}| = |\mathcal{T}_1| + |\mathcal{T}_2| \leq (\frac{k}{2} - 1) + (n - k - 1) = n - \frac{k}{2} - 2 < n - \frac{k}{2} - 1$, as desired.

303 Consider the case $|\mathcal{T}_1| = \frac{k}{2}$. We claim that $|\mathcal{T}_2| \leq n - k - 2$. Assume, to the
 304 contrary, that $n - k - 1 \leq |\mathcal{T}_2| \leq n - k$. Since $|\mathcal{T}_1| = \frac{k}{2}$, it follows that each edge
 305 of $G[S]$ is occupied by some tree in \mathcal{T}_1 , which implies that each tree in \mathcal{T}_2 only
 306 uses the edges of $E_G[S, \bar{S}] \cup E(G[\bar{S}])$. Suppose that T_1 is a tree in \mathcal{T}_2 occupying
 307 w_1 . Since $u_1 w_1 \notin E(G)$, if T_1 contains three vertices of \bar{S} , then the remaining
 308 $n - k - 3$ vertices in \bar{S} must be contained in at most $n - k - 3$ trees in \mathcal{T}_2 , which
 309 results in $|\mathcal{T}_2| \leq (n - k - 3) + 1 = n - k - 2$, a contradiction. So we assume that the
 310 tree T_1 contains another vertex of \bar{S} except w_1 , say w_2 . Recall that $k \geq 4$. Then
 311 $|\bar{S}| \geq k \geq 4$. By the same reason, there is another tree T_2 containing two vertices
 312 of \bar{S} , say w_3, w_4 . Furthermore, the remaining $n - k - 4$ vertices in \bar{S} must be
 313 contained in at most $n - k - 4$ trees in \mathcal{T}_2 , which results in $|\mathcal{T}_2| \leq (n - k - 4) + 2 =$
 314 $n - k - 2$, a contradiction. We conclude that $|\mathcal{T}_2| \leq n - k - 2$. Since $|\mathcal{T}_1| = \frac{k}{2}$, we
 315 have $\kappa(S) \leq \lambda(S) = |\mathcal{T}| = |\mathcal{T}_1| + |\mathcal{T}_2| \leq \frac{k}{2} + (n - k - 2) = n - \frac{k}{2} - 2 < n - \frac{k}{2} - 1$,
 316 as desired. ■

317 **Lemma 2.6.** *If n ($n \geq 4$) is even and M is a set of edges in the complete graph*
 318 *K_n such that $1 \leq |M| \leq n - 1$ and $1 \leq \Delta(K_n[M]) \leq \frac{n}{2}$, then $G = K_n - M$*
 319 *contains $\frac{n-2}{2}$ edge-disjoint spanning trees.*

320 **Proof.** Let $\mathcal{P} = \bigcup_{i=1}^p V_i$ be a partition of $V(G)$ with $|V_i| = n_i$ ($1 \leq i \leq p$), and
 321 \mathcal{E}_p be the set of edges between distinct blocks of \mathcal{P} in G . It suffices to show that
 322 $|\mathcal{E}_p| \geq \frac{n-2}{2}(|\mathcal{P}| - 1)$ so that we can use Theorem 1.1.

323 The case $p = 1$ is trivial by Theorem 1.1, thus we assume $p \geq 2$. For
 324 $p = 2$, we have $\mathcal{P} = V_1 \cup V_2$. Set $|V_1| = n_1$. Clearly, $|V_2| = n - n_1$. Since
 325 $\Delta(K_n[M]) \leq \frac{n}{2}$, it follows that $\delta(G) = n - 1 - \Delta(K_n[M]) \geq n - 1 - \frac{n}{2} = \frac{n-2}{2}$.
 326 Therefore, if $n_1 = 1$ then $|\mathcal{E}_2| = |E_G[V_1, V_2]| \geq \frac{n-2}{2}$. Suppose $n_1 \geq 2$. Then
 327 $|\mathcal{E}_2| = |E_G[V_1, V_2]| \geq \binom{n}{2} - (n-1) - \binom{n_1}{2} - \binom{n-n_1}{2} = -n_1^2 + nn_1 - n + 1$. Since
 328 $2 \leq n_1 \leq n - 2$, one can see that $|\mathcal{E}_2|$ achieves its minimum value when $n_1 = 2$
 329 or $n_1 = n - 2$. Thus $|\mathcal{E}_2| \geq n - 3 \geq \frac{n-2}{2}$ since $n \geq 4$. The result follows from
 330 Theorem 1.1.

331 Let us consider the remaining cases for p , namely, for $3 \leq p \leq n$. Since
 332 $|\mathcal{E}_p| \geq \binom{n}{2} - |M| - \sum_{i=1}^p \binom{n_i}{2} \geq \binom{n}{2} - (n-1) - \sum_{i=1}^p \binom{n_i}{2} = \binom{n-1}{2} - \sum_{i=1}^p \binom{n_i}{2}$, we
 333 only need to show $\binom{n-1}{2} - \sum_{i=1}^p \binom{n_i}{2} \geq \frac{n-2}{2}(p-1)$, that is, $(n-p)\frac{n-2}{2} \geq \sum_{i=1}^p \binom{n_i}{2}$.
 334 Because $\sum_{i=1}^p \binom{n_i}{2}$ achieves its maximum value when $n_1 = n_2 = \dots = n_{p-1} = 1$

335 and $n_p = n - p + 1$, we need inequality $(n - p)\frac{n-2}{2} \geq \binom{1}{2}(p-1) + \binom{n-p+1}{2}$, namely,
 336 $(n - p)\frac{n-3}{2} \geq 0$. It is easy to see that the inequality holds since $3 \leq p \leq n$. Thus,
 337 $|\mathcal{E}_p| \geq \binom{n}{2} - |M| - \sum_{i=1}^p \binom{n_i}{2} \geq \frac{n-2}{2}(p-1)$.

338 From Theorem 1.1, there exist $\frac{n-2}{2}$ edge-disjoint spanning trees in G , as
 339 desired. \blacksquare

340 **Lemma 2.7.** *Let k, n be two integers with $4 \leq k \leq n$, and M is an edge set of*
 341 *the complete graph K_n satisfying $\Delta(K_n[M]) = 1$. Then*

342 (1) *If $|M| = k - 1$, then $\kappa_k(K_n - M) \geq n - \frac{k}{2} - 1$;*

343 (2) *If $|M| = \lfloor \frac{n}{2} \rfloor$, then $\lambda_k(K_n - M) \geq n - \frac{k}{2} - 1$.*

344 **Proof.** (1) Set $G = K_n - M$. Since $\Delta(K_n[M]) = 1$, it follows that M is a
 345 matching of K_n . By the definition of $\kappa_k(G)$, we need to show that $\kappa(S) \geq n - \frac{k}{2} - 1$
 346 for any $S \subseteq V(G)$.

347 **Case 1.** There exists no u, w in S such that $uw \in M$.

348 Without loss of generality, let $S = \{u_1, u_2, \dots, u_k\}$ such that u_1, u_2, \dots, u_r
 349 are M -saturated but $u_{r+1}, u_{r+2}, \dots, u_k$ are M -unsaturated. Let $M_1 = \{u_i w_i \mid 1 \leq$
 350 $i \leq r\} \subseteq M$. Since $|M| = k - 1$, it follows that $0 \leq r \leq k - 1$. In this case,
 351 $u_i u_j \notin M$ ($1 \leq i, j \leq r$). Clearly, $G[S]$ is a clique of order k . We choose a path
 352 $P = u_1 u_2 \dots u_r u_{r+1}$ in $G[S]$. Let $G' = G - E(P)$. Then $G'[S] = K_k - E(P)$.
 353 Since $|E(P)| = r \leq k - 1$ and $\Delta(K_k[E(P)]) = 2 \leq \frac{k}{2}$, it follows that $G'[S]$
 354 contains $\frac{k-2}{2}$ edge-disjoint spanning trees, which are also $\frac{k-2}{2}$ internally disjoint
 355 S -Steiner trees. These trees together with the trees T_i induced by the edges
 356 in $\{u_1 w_i, u_2 w_i, u_{i-1} w_i, u_{i+1} w_i, \dots, u_k w_i, u_i u_{i+1}\}$ ($1 \leq i \leq r$) (see Figure 1 (a))
 357 and the trees T_j induced by the edges in $\{u_1 v_j, u_2 v_j, \dots, u_k v_j\}$ where $v_j \in \bar{S} -$
 358 $\{w_1, w_2, \dots, w_r\} = \{v_1, v_2, \dots, v_{n-k-r}\}$ form $\frac{k-2}{2} + r + (n - k - r) = n - \frac{k}{2} - 1$
 359 internally disjoint S -Steiner trees. Thus, $\kappa(S) \geq n - \frac{k}{2} - 1$, as desired.

360 **Case 2.** There exist u, w in S such that $uw \in M$.

361 Without loss of generality, we let $S = \{u_1, u_2, \dots, u_r, u_{r+1}, u_{r+2}, \dots, u_{r+s},$
 362 $u_{r+s+1}, \dots, u_{k-r}, w_1, w_2, \dots, w_r\}$ such that the vertices $u_1, u_2, \dots, u_{r+s}, w_1, w_2,$
 363 \dots, w_r are all M -saturated and $u_i w_i \in M$ ($1 \leq i \leq r$). Set $M_1 = \{u_i w_i \mid 1 \leq$
 364 $i \leq r\}$. In this case, $r \geq 1$ and $2r + s \leq k$. Since $|M| = k - 1$, it follows that
 365 $r + s \leq k - 1$ and $s \leq k - 2$.

366 First, we consider $2r + s = k$. Since k is even, it follows that s is even.
 367 If $s = 0$, then $r = \frac{k}{2}$. Thus $S = \{u_1, u_2, \dots, u_{\frac{k}{2}}, w_1, w_2, \dots, w_{\frac{k}{2}}\}$. Clearly,
 368 $M_1 = \{u_i w_i \mid 1 \leq i \leq \frac{k}{2}\}$, $|M_1| = \frac{k}{2} \leq k - 1$ and $\Delta(K_n[M_1]) = 1 < \frac{k}{2}$. By
 369 Lemma 2.6, $G[S]$ contains $\frac{k-2}{2}$ edge-disjoint spanning trees, which are also $\frac{k-2}{2}$
 370 internally disjoint S -Steiner trees. These trees together with the trees T_j induced
 371 by the edges in $\{u_1 v_j, u_2 v_j, \dots, u_{\frac{k}{2}} v_j\} \cup \{w_1 v_j, w_2 v_j, \dots, w_{\frac{k}{2}} v_j\}$ form $\frac{k-2}{2} + (n - k)$
 372 internally disjoint S -Steiner trees, where $v_j \in \bar{S} = \{v_1, v_2, \dots, v_{n-k}\}$. So, $\kappa(S) \geq$
 373 $n - \frac{k}{2} - 1$.




Figure 1. Graphs for (1) of Lemma 2.7.

374 Consider $s = 2$. Since $2r + s = k$, we have $r = \frac{k-2}{2}$. If $k = 4$, then
 375 $r = 1$ and hence $S = \{u_1, u_2, u_3, w_1\}$. Clearly, $M_1 = \{u_1 w_1\}$, and the tree
 376 T_1 induced by the edges in $\{u_1 u_2, u_1 w_2, w_1 w_2, u_3 w_2\}$ and the tree T_2 induced
 377 by the edges in $\{u_1 u_3, u_2 u_3, u_2 w_1\}$ and the tree T_3 induced by the edges in
 378 $\{u_1 w_3, u_2 w_3, w_1 w_3, u_3 w_1\}$ are three spanning trees; see Figure 1 (c). These trees
 379 together with the trees T_j induced by the edges in $\{u_1 v_j, u_2 v_j, u_3 v_j, w_1 v_j\}$ form
 380 $3 + (n - 6)$ internally disjoint S -Steiner trees, where $v_j \in \bar{S} - \{w_2, w_3\} =$
 381 $\{v_1, v_2, \dots, v_{n-6}\}$. Thus, $\kappa(S) \geq n - 3 = n - \frac{k}{2} - 1$. Suppose $k \geq 6$. Then
 382 $r \geq 2$, $S = \{u_1, u_2, \dots, u_{\frac{k+2}{2}}, w_1, w_2, \dots, w_{\frac{k-2}{2}}\}$ and $M_1 = \{u_i w_i \mid 1 \leq i \leq \frac{k-2}{2}\}$.
 383 Clearly, the tree T_1 induced by the edges in $\{u_1 w_{\frac{k}{2}}, u_2 w_{\frac{k}{2}}, \dots, u_{\frac{k-2}{2}} w_{\frac{k}{2}}, u_{\frac{k+2}{2}} w_{\frac{k}{2}},$
 384 $u_2 u_{\frac{k}{2}}, w_1 w_{\frac{k}{2}}, w_2 w_{\frac{k}{2}}, \dots, w_{\frac{k-2}{2}} w_{\frac{k}{2}}\}$ and the tree T_2 induced by the edges in $\{u_1 w_{\frac{k+2}{2}},$
 385 $u_2 w_{\frac{k+2}{2}}, \dots, u_{\frac{k}{2}} w_{\frac{k+2}{2}}\} \cup \{u_1 u_{\frac{k+2}{2}}, w_1 w_{\frac{k+2}{2}}, w_2 w_{\frac{k+2}{2}}, \dots, w_{\frac{k-2}{2}} w_{\frac{k+2}{2}}\}$ are two inter-
 386 nally disjoint S -Steiner trees; see Figure 1 (d). Let $M_2 = M_1 \cup \{u_1 u_{\frac{k+2}{2}}, u_2 u_{\frac{k}{2}}\}$.
 387 Then $|M_2| = |M_1| + 2 = \frac{k-2}{2} + 2 = \frac{k+2}{2} < k - 1$ and $\Delta(K_n[M_2]) = 2 \leq \frac{k}{2}$,
 388 which implies that $G[S] - \{u_1 u_{\frac{k+2}{2}}, u_2 u_{\frac{k}{2}}\} = K_k - M_2$ contains $\frac{k-2}{2}$ edge-disjoint
 389 spanning trees by Lemma 2.6, which are also $\frac{k-2}{2}$ internally disjoint S -Steiner
 390 trees. These trees together with T_1, T_2 and the trees T_j induced by the edges in
 391 $\{u_1 v_j, u_2 v_j, \dots, u_{\frac{k+2}{2}} v_j, w_1 v_j, w_2 v_j, \dots, u_{\frac{k-2}{2}} v_j\}$ are $\frac{k-2}{2} + 2 + (n - k - 2)$ inter-
 392 nally disjoint S -Steiner trees, where $v_j \in \bar{S} - \{w_{\frac{k}{2}}, w_{\frac{k+2}{2}}\} = \{v_1, v_2, \dots, v_{n-k-2}\}$.
 393 So, $\kappa(S) \geq n - \frac{k}{2} - 1$.

394 Consider the remaining case for s , namely, for $4 \leq s \leq k - 2$. Clearly,
 395 there exists a cycle of order s containing $u_{r+1}, u_{r+2}, \dots, u_{r+s}$ in $K_k - M_1$, say

396 $C_s = u_{r+1}u_{r+2} \cdots u_{r+s}u_{r+1}$. Set $M' = M_1 \cup E(C_s)$. Then $|M'| = r + s \leq k - 1$
 397 and $\Delta(K_n[M']) = 2 \leq \frac{k}{2}$, which implies that $G - E(C_s) = K_k - M'$ contains $\frac{k-2}{2}$
 398 edge-disjoint spanning trees by Lemma 2.6. These trees together with the trees
 399 T_{r+j} induced by the edges in $\{u_1w_{r+j}, u_2w_{r+j}, \dots, u_{r+j-1}w_{r+j}, u_{r+j+1}w_{r+j}, \dots,$
 400 $u_{r+s}w_{r+j}, u_{r+j}u_{r+j+1}, w_1w_{r+j}, w_2w_{r+j}, \dots, w_rw_{r+j}\}$ ($1 \leq j \leq s$) form $\frac{k-2}{2} + s$
 401 internally disjoint trees; see Figure 2 (b) (note that $u_{r+s} = u_{k-r}$). These trees to-
 402 gether with the trees T'_j induced by the edges in $\{u_1v_j, u_2v_j, \dots, u_{r+s}v_j, w_1v_j, \dots,$
 403 $w_rv_j\}$ form $\frac{k-2}{2} + s + (n - 2r - 2s) = n - \frac{k}{2} - 1$ internally disjoint S -Steiner
 404 trees where $v_j \in \bar{S} - \{w_{r+1}, w_{r+2}, \dots, w_{r+s}\} = \{v_1, v_2, \dots, v_{n-2r-2s}\}$. Thus,
 405 $\kappa(S) \geq n - \frac{k}{2} - 1$, as desired.

406 Next, assume $2r + s < k$. Then $S = \{u_1, u_2, \dots, u_{r+s}, u_{r+s+1}, \dots, u_{k-r}, w_1,$
 407 $w_2, \dots, w_r\}$ and $r+s+1 \leq k-r$. If $s = 0$, then $S = \{u_1, u_2, \dots, u_{k-r}, w_1, w_2, \dots,$
 408 $w_r\}$. Clearly, $M_1 = \{u_iw_i \mid 1 \leq i \leq r\}$, $|M_1| = r \leq k-1$ and $\Delta(K_n[M_1]) = 1 < \frac{k}{2}$.
 409 By Lemma 2.6, $G[S]$ contains $\frac{k-2}{2}$ edge-disjoint spanning trees. These trees to-
 410 gether with the trees T_j induced by the edges in $\{u_1v_j, u_2v_j, \dots, u_{n-r}v_j, w_1v_j, w_2v_j,$
 411 $\dots, w_rv_j\}$ form $\frac{k-2}{2} + (n - k)$ internally disjoint S -Steiner trees, where $v_j \in \bar{S} =$
 412 $\{v_1, v_2, \dots, v_{n-k}\}$. Therefore, $\kappa(S) \geq n - \frac{k}{2} - 1$. Assume $s \geq 1$. Clearly, there
 413 exists a path of length s containing $u_{r+1}, u_{r+2}, \dots, u_{r+s}, u_{r+s+1}$ in $G[S]$, say
 414 $P_s = u_{r+1}u_{r+2} \cdots u_{r+s}u_{r+s+1}$. Set $M' = M_1 \cup E(P_s)$. Then $|M'| = r + s \leq k - 1$
 415 and $\Delta(K_n[M']) = 2 \leq \frac{k}{2}$, which implies that $G[S] - E(P_s) = K_k - M'$ contains $\frac{k-2}{2}$
 416 edge-disjoint spanning trees by Lemma 2.6, which are also $\frac{k-2}{2}$ internally disjoint
 417 S -Steiner trees. These trees together with the trees T_{r+j} induced by the edges in
 418 $\{u_1w_{r+j}, u_2w_{r+j}, \dots, u_{r+j-1}w_{r+j}, u_{r+j+1}w_{r+j}, \dots, u_{k-r}w_{r+j}, u_{r+j}u_{r+j+1}, w_1w_{r+j},$
 419 $w_2w_{r+j}, \dots, w_rw_{r+j}\}$ ($1 \leq j \leq s$) form $\frac{k-2}{2} + s$ internally disjoint S -Steiner
 420 trees; see Figure 1 (b). These trees together with the trees T'_j induced by
 421 the edges in $\{u_1v_j, u_2v_j, \dots, u_{k-r}v_j, w_1v_j, w_2v_j, \dots, w_rv_j\}$ form $\frac{k-2}{2} + s + (n -$
 422 $k + r) - (r + s) = n - \frac{k}{2} - 1$ internally disjoint S -Steiner trees where $v_j \in$
 423 $\bar{S} - \{w_{r+1}, w_{r+2}, \dots, w_{r+s}\} = \{v_1, v_2, \dots, v_{n-k-s}\}$. So, $\kappa(S) \geq n - \frac{k}{2} - 1$, as
 424 desired.

425 We conclude that $\kappa(S) \geq n - \frac{k}{2} - 1$ for any $S \subseteq V(G)$. From the arbitrariness
 426 of S , it follows that $\kappa_k(G) \geq n - \frac{k}{2} - 1$.

427 (2) Set $G = K_n - M$. Assume that n is even. Thus M is a perfect matching
 428 of K_n , and all vertices of G are M -saturated. By the definition of $\lambda_k(G)$, we need
 429 to show that $\lambda(S) \geq n - \frac{k}{2} - 1$ for any $S \subseteq V(G)$.

430 **Case 3.** There exists no u, w in S such that $uw \in M$.

431 Without loss of generality, let $S = \{u_1, u_2, \dots, u_k\}$. In this case, $u_iu_j \notin$
 432 M ($1 \leq i, j \leq k$). Let $M_1 = \{u_iw_i \mid 1 \leq i \leq k\} \subseteq M = \{u_iw_i \mid 1 \leq i \leq \frac{n}{2}\}$.
 433 Clearly, $w_i \notin S$ ($1 \leq i \leq \frac{n}{2}$) and $u_j \notin S$ ($k+1 \leq j \leq \frac{n}{2}$). Since $G[S]$ is a clique of
 434 order k , it follows that there are $\frac{k}{2}$ edge-disjoint spanning trees in $G[S]$, which are
 435 also $\frac{k}{2}$ edge-disjoint S -Steiner trees. These trees together with the trees T_i induced

436 by the edges in $\{u_1w_i, u_2w_i, u_{i-1}w_i, u_{i+1}w_i, \dots, u_kw_i, u_iw_k, w_iw_k\}$ ($1 \leq i \leq k-1$)
 437 (see Figure 2 (a)) and the trees T'_j induced by the edges in $\{u_1u_j, u_2u_j, \dots, u_ku_j\}$
 438 ($k+1 \leq j \leq \frac{n}{2}$) and the trees T''_j induced by the edges in $\{u_1w_j, u_2w_j, \dots, u_kw_j\}$
 439 ($k+1 \leq j \leq \frac{n}{2}$) form $\frac{k}{2} + (k-1) + (n-2k) = n - \frac{k}{2} - 1$ edge-disjoint S -Steiner
 440 trees. Therefore, $\lambda(S) \geq n - \frac{k}{2} - 1$, as desired.

441 **Case 4.** There exist u, w in S such that $uw \in M$.

442 Without loss of generality, let $S = \{u_1, u_2, \dots, u_{r+s}, w_1, w_2, \dots, w_r\}$ with
 443 $|S| = k = 2r + s$, where $1 \leq r \leq \frac{k}{2}$ and $0 \leq s \leq k - 2$. Set $M_1 = \{u_iw_i \mid 1 \leq$
 444 $i \leq r\} \subseteq M = \{u_iw_i \mid 1 \leq i \leq \frac{n}{2}\}$. We claim that $r + s \leq k - 1$. Otherwise, let
 445 $r + s = k$. Combining this with $2r + s = k$, we have $r = 0$, a contradiction. Since
 $k = 2r + s$ and k is even, it follows that s is even.




Figure 2. Graphs for (2) of Lemma 2.7.

446 If $s = 0$, then $r = \frac{k}{2}$. Clearly, $S = \{u_1, u_2, \dots, u_{\frac{k}{2}}, w_1, w_2, \dots, w_{\frac{k}{2}}\}$ and
 447 $M_1 = M = \{u_iw_i \mid 1 \leq i \leq \frac{k}{2}\}$. In addition, $|M_1| \leq \frac{k}{2} < k - 1$ and $\Delta(M \cap$
 448 $K_n[S]) = 1 < \frac{k}{2}$. Then $G[S]$ contains $\frac{k-2}{2}$ edge-disjoint spanning trees by
 449 Lemma 2.6. These trees together with the trees T_i induced by the edges in
 450 $\{u_1u_i, u_2u_i, \dots, u_{\frac{k}{2}}u_i, w_1u_i, w_2u_i, \dots, w_{\frac{k}{2}}u_i\}$ ($k+1 \leq j \leq \frac{n}{2}$) and the trees T'_i
 451 induced by the edges in $\{u_1w_i, u_2w_i, \dots, u_{\frac{k}{2}}w_i, w_1w_i, w_2w_i, \dots, w_{\frac{k}{2}}w_i\}$ ($\frac{k}{2} + 1 \leq$
 452 $i \leq \frac{n}{2}$) form $n - \frac{k}{2} - 1$ edge-disjoint S -Steiner trees. Thus, $\lambda(S) \geq n - \frac{k}{2} - 1$.
 453

454 If $s = 2$, then $r = \frac{k-2}{2}$. Then $S = \{u_1, u_2, \dots, u_{\frac{k+2}{2}}, w_1, w_2, \dots, w_{\frac{k-2}{2}}\}$
 455 and $M_1 = \{u_iw_i \mid 1 \leq i \leq \frac{k-2}{2}\} \subseteq M$. If $k = 4$, then $r = 1$ and hence $S =$
 456 $\{u_1, u_2, u_3, w_1\}$. Clearly, $M_1 = \{u_1w_1\}$, and the tree T_1 induced by the edges in
 457 $\{u_1u_2, u_1w_2, w_1w_2, u_3w_2\}$ and the tree T_2 induced by the edges in $\{u_1u_3, u_2u_3, u_2w_1\}$
 458 and the tree T_3 induced by the edges in $\{u_1w_3, u_2w_3, w_1w_3, u_3w_1\}$ are three edge-
 459 disjoint spanning trees; see Figure 1 (c). These trees together with the trees T_j
 460 induced by the edges in $\{u_1u_j, u_2u_j, u_3u_j, w_1u_j\}$ ($4 \leq k \leq \frac{n}{2}$) and the trees T'_j in-
 461 duced by the edges in $\{u_1w_j, u_2w_j, u_3w_j, w_1w_j\}$ ($4 \leq k \leq \frac{n}{2}$) form $3 + (n-6)$ edge-
 462 disjoint S -Steiner trees. So, $\lambda(S) \geq n - 3 = n - \frac{k}{2} - 1$, as desired. Suppose $k \geq 6$.
 463 Then $r \geq 2$, $S = \{u_1, u_2, \dots, u_{\frac{k+2}{2}}, w_1, w_2, \dots, w_{\frac{k-2}{2}}\}$ and $M_1 = \{u_iw_i \mid 1 \leq i \leq$

464 $\frac{k-2}{2}$. Clearly, the tree T_1 induced by the edges in $\{u_1w_{\frac{k}{2}}, u_2w_{\frac{k}{2}}, \dots, u_{\frac{k-2}{2}}w_{\frac{k}{2}},$
 465 $u_{\frac{k+2}{2}}w_{\frac{k}{2}}, u_2u_{\frac{k}{2}}, w_1w_{\frac{k}{2}}, w_2w_{\frac{k}{2}}, \dots, w_{\frac{k-2}{2}}w_{\frac{k}{2}}\}$ and the tree T_2 induced by the edges
 466 in $\{u_1w_{\frac{k+2}{2}}, u_2w_{\frac{k+2}{2}}, \dots, u_{\frac{k}{2}}w_{\frac{k+2}{2}}, u_1u_{\frac{k+2}{2}}, w_1w_{\frac{k+2}{2}}, w_2w_{\frac{k+2}{2}}, \dots, w_{\frac{k-2}{2}}w_{\frac{k+2}{2}}\}$ are
 467 two edge-disjoint S -Steiner trees; see Figure 1 (d). Let $M_2 = M_1 \cup \{u_1u_{\frac{k+2}{2}}, u_2u_{\frac{k}{2}}\}$.
 468 Then $|M_2| = |M_1| + 2 = \frac{k-2}{2} + 2 = \frac{k+2}{2} < k-1$ and $\Delta(K_n[M_2]) = 2 \leq \frac{k}{2}$, which im-
 469 plies that $G[S] - \{u_1u_{\frac{k+2}{2}}, u_2u_{\frac{k}{2}}\} = K_k - M_2$ contains $\frac{k-2}{2}$ edge-disjoint spanning
 470 trees by Lemma 2.6. These trees together with T_1, T_2 and the trees T_j induced by
 471 the edges in $\{u_1u_j, u_2u_j, \dots, u_{\frac{k+2}{2}}u_j, w_1u_j, w_2u_j, \dots, u_{\frac{k-2}{2}}u_j\}$ ($\frac{k}{2} + 2 \leq j \leq \frac{n}{2}$)
 472 and the trees T'_j induced by the edges in $\{u_1w_j, u_2w_j, \dots, u_{\frac{k+2}{2}}w_j, w_1w_j, w_2w_j,$
 473 $\dots, u_{\frac{k-2}{2}}w_j\}$ ($\frac{k}{2} + 2 \leq j \leq \frac{n}{2}$) are $\frac{k-2}{2} + 2 + (n-k-2)$ edge-disjoint S -Steiner
 474 trees. Therefore, $\lambda(S) \geq n - \frac{k}{2} - 1$, as desired.

475 Consider the remaining case s with $4 \leq s \leq k-2$. Clearly, there ex-
 476 ists a cycle of order s containing $u_{r+1}, u_{r+2}, \dots, u_{r+s}$ in $K_k - M_1$, say $C_s =$
 477 $u_{r+1}u_{r+2} \dots u_{r+s}u_{r+1}$. Set $M' = M_1 \cup E(C_s)$. Then $|M'| = r + s \leq k-1$ and
 478 $\Delta(K_n[M']) = 2 \leq \frac{k}{2}$, which implies that $G - E(C_s)$ contains $\frac{k-2}{2}$ edge-disjoint s-
 479 panning trees by Lemma 2.6. These trees together with the trees T_{r+j} induced by
 480 the edges in $\{u_1w_{r+j}, u_2w_{r+j}, \dots, u_{r+j-1}w_{r+j}, u_{r+j+1}w_{r+j}, \dots, u_{r+s}w_{r+j}, u_{r+j}$
 481 $u_{r+j+1}, w_1w_{r+j}, w_2w_{r+j}, \dots, w_rw_{r+j}\}$ ($1 \leq j \leq s$) form $\frac{k-2}{2} + s$ edge-disjoint S -
 482 Steiner trees; see Figure 2 (b). These trees together with the trees T'_i induced by
 483 the edges in $\{u_1u_i, u_2u_i, \dots, u_{r+s}u_i, w_1u_i, \dots, w_ru_i\}$ ($r+s+1 \leq i \leq \frac{n}{2}$) and the
 484 trees T''_i induced by the edges in $\{u_1w_i, u_2w_i, \dots, u_{r+s}w_i, w_1w_i, \dots, w_rw_i\}$ ($r+$
 485 $s+1 \leq i \leq \frac{n}{2}$) form $(n-2r-2s) + (\frac{k-2}{2} + s) = n - \frac{k}{2} - 1$ edge-disjoint S -Steiner
 486 trees since $2r+s = k$. Thus, $\lambda(S) \geq n - \frac{k}{2} - 1$, as desired.

487 We conclude that $\lambda(S) \geq n - \frac{k}{2} - 1$ for any $S \subseteq V(G)$. From the arbitrariness
 488 of S , it follows that $\lambda_k(G) \geq n - \frac{k}{2} - 1$. For n odd, M is a maximum matching
 489 and we can also check that $\lambda_k(G) \geq n - \frac{k}{2} - 1$ similarly. \blacksquare

490 **Lemma 2.8.** *Let n and k be two integers such that k is even and $4 \leq k \leq n$.
 491 If M is a set of edges in the complete graph K_n such that $|M| = k-1$, and
 492 $2 \leq \Delta(K_n[M]) \leq \frac{k}{2}$, then $\kappa_k(K_n - M) \geq n - \frac{k}{2} - 1$.*

493 **Proof.** Set $G = K_n - M$. For $n = k$, there are $\frac{n-2}{2}$ edge-disjoint spanning trees
 494 by Lemma 2.6, and hence $\kappa_n(G) = \lambda_n(G) \geq \frac{n-2}{2}$. So from now on, we assume $n \geq$
 495 $k+1$. Let $S = \{u_1, u_2, \dots, u_k\} \subseteq V(G)$ and $\bar{S} = V(G) - S = \{w_1, w_2, \dots, w_{n-k}\}$.
 496 We have the following two cases to consider.

497 **Case 1.** $M \subseteq E(K_n[S]) \cup E(K_n[\bar{S}])$.

498 Let $M' = M \cap E(K_n[S])$ and $M'' = M \cap E(K_n[\bar{S}])$. Then $|M'| + |M''| =$
 499 $|M| = k-1$ and $0 \leq |M'|, |M''| \leq k-1$. We can regard $G[S]$ as a complete
 500 graph K_k by deleting $|M'|$ edges. Since $2 \leq \Delta(K_n[M]) \leq \frac{k}{2}$ and $M' \subseteq M$, it
 501 follows that $\Delta(K_n[M']) \leq \Delta(K_n[M]) \leq \frac{k}{2}$. From Lemma ??, there exist $\frac{k-2}{2}$

502 edge-disjoint spanning trees in $G[S]$. Actually, these $\frac{k-2}{2}$ edge-disjoint spanning
 503 trees are all internally disjoint S -Steiner trees in $G[S]$. All these trees together
 504 with the trees T_i induced by the edges in $\{w_i u_1, w_i u_2, \dots, w_i u_k\}$ ($1 \leq i \leq n-k$)
 505 form $\frac{k-2}{2} + (n-k) = n - \frac{k}{2} - 1$ internally disjoint S -Steiner trees, and hence
 506 $\kappa(S) \geq n - \frac{k}{2} - 1$. From the arbitrariness of S , we have $\kappa_k(G) \geq n - \frac{k}{2} - 1$, as
 507 desired.

508 **Case 2.** $M \not\subseteq E(K_n[S]) \cup E(K_n[\bar{S}])$.

509 In this case, there exist some edges of M in $E_{K_n}[S, \bar{S}]$. Let $M' = M \cap$
 510 $E(K_n[S])$, $M'' = M \cap E(K_n[\bar{S}])$, and $|M'| = m_1$ and $|M''| = m_2$. Clearly, $0 \leq$
 511 $m_i \leq k-2$ ($i = 1, 2$). For $w_i \in \bar{S}$, let $|E_{K_n[M]}[w_i, S]| = x_i$, where $1 \leq i \leq n-k$.
 512 Without loss of generality, let $x_1 \geq x_2 \geq \dots \geq x_{n-k}$. Because there exist some
 513 edges of M in $E_{K_n}[S, \bar{S}]$, we have $x_1 \geq 1$. Since $2 \leq \Delta(K_n[M]) \leq \frac{k}{2}$, it follows
 514 that $x_i = |E_{K_n[M]}[w_i, S]| \leq d_{K_n[M]}(w_i) \leq \Delta(K_n[M]) \leq \frac{k}{2}$ for $1 \leq i \leq n-k$.
 515 We claim that there exists at most one vertex in $K_n[M]$ such that its degree is
 516 $\frac{k}{2}$. Assume, to the contrary, that there are two vertices, say w and w' , such that
 517 $d_{K_n[M]}(w) = d_{K_n[M]}(w') = \frac{k}{2}$. Then $|M| \geq d_{K_n[M]}(w) + d_{K_n[M]}(w') = \frac{k}{2} + \frac{k}{2} = k$,
 518 contradicting $|M| = k-1$. We conclude that there exists at most one vertex in
 519 $K_n[M]$ such that its degree is $\frac{k}{2}$. Recall that $x_{n-k} \leq x_{n-k-1} \leq \dots \leq x_2 \leq x_1 \leq \frac{k}{2}$.
 520 So $x_1 = \frac{k}{2}$ and $x_i \leq \frac{k-2}{2}$ ($2 \leq i \leq n-k$), or $x_i \leq \frac{k-2}{2}$ ($1 \leq i \leq n-k$). Since
 521 $|E_{K_n[M]}[w_i, S]| = x_i$, we have $|E_G[w_i, S]| = k - x_i$. Since $2 \leq \Delta(K_n[M]) \leq \frac{k}{2}$, it
 522 follows that $\delta(G[S]) \geq k-1 - \frac{k}{2} = \frac{k-2}{2}$.

523 Our basic idea is to seek for some edges in $G[S]$, and combine them with
 524 the edges of $E_G[S, \bar{S}]$ to form $n-k$ internally disjoint trees, say T_1, T_2, \dots, T_{n-k} ,
 525 with their roots w_1, w_2, \dots, w_{n-k} , respectively. Let $G' = G - (\bigcup_{j=1}^{n-k} E(T_j))$. We
 526 will prove that $G'[S]$ satisfies the conditions of Lemma ?? so that $G'[S]$ contains
 527 $\frac{k-2}{2}$ edge-disjoint spanning trees, which are also $\frac{k-2}{2}$ internally disjoint S -Steiner
 528 trees. These trees together with T_1, T_2, \dots, T_{n-k} are our $n - \frac{k}{2} - 1$ desired trees.
 529 Thus, $\kappa(S) \geq n - \frac{k}{2} - 1$. So we can complete our proof by the arbitrariness of S .

530 For $w_1 \in \bar{S}$, without loss of generality, let $S = S_1^1 \cup S_2^1$ and $S_1^1 = \{u_1, u_2, \dots,$
 531 $u_{x_1}\}$ such that $u_j w_1 \in M$ for $1 \leq j \leq x_1$. Set $S_2^1 = S - S_1^1 = \{u_{x_1+1}, u_{x_1+2}, \dots, u_k\}$.
 532 Then $u_j w_1 \in E(G)$ for $x_1 + 1 \leq j \leq k$. One can see that the tree T_1' induced
 533 by the edges in $\{w_1 u_{x_1+1}, w_1 u_{x_1+2}, \dots, w_1 u_k\}$ is a Steiner tree connecting S_2^1 .
 534 Our current idea is to seek for x_1 edges in $E_G[S_1^1, S_2^1]$ and add them to T_1' to
 535 form a Steiner tree connecting S . For each $u_j \in S_1^1$ ($1 \leq j \leq x_1$), we claim that
 536 $|E_G[u_j, S_2^1]| \geq 1$. Otherwise, let $|E_G[u_j, S_2^1]| = 0$. Then $|E_{K_n[M]}[u_j, S_2^1]| = k - x_1$
 537 and hence $|M| \geq |E_{K_n[M]}[u_j, S_2^1]| + d_{K_n[M]}(w_1) \geq (k - x_1) + x_1 = k$, which con-
 538 tradicts $|M| = k-1$. We conclude that for each $u_j \in S_1^1$ ($1 \leq j \leq x_1$) there is
 539 at least one edge in G connecting it to a vertex of S_2^1 . Choose the vertex with
 540 the smallest subscript among all the vertices of S_2^1 having maximum degree in
 541 $G[S]$, say u_1' . Then we select the vertex adjacent to u_1' with the smallest sub-

542 script among all the vertices of S_2^1 having maximum degree in $G[S]$, say u_1'' . Let
 543 $e_{11} = u_1' u_1''$. Consider the graph $G_{11} = G - e_{11}$, and choose the vertex with
 544 the smallest subscript among all the vertices of $S_1^1 - u_1'$ having maximum degree
 545 in $G_{11}[S]$, say u_2' . Then we select the vertex adjacent to u_2' with the smallest
 546 subscript among all the vertices of S_2^1 having maximum degree in $G_{11}[S]$, say u_2'' .
 547 Set $e_{12} = u_2' u_2''$. Consider the graph $G_{12} = G_{11} - e_{12} = G - \{e_{11}, e_{12}\}$. Choose
 548 the one with the smallest subscript among all the vertices of $S_1^1 - \{u_1', u_2'\}$ having
 549 maximum degree in $G_{12}[S]$, say u_3' , and select the vertex adjacent to u_3' with the
 550 smallest subscript among all the vertices of S_2^1 having maximum degree in $G_{12}[S]$,
 551 say u_3'' . Put $e_{13} = u_3' u_3''$. Consider the graph $G_{13} = G_{12} - e_{11} = G - \{e_{11}, e_{12}, e_{13}\}$.
 552 For each $u_j \in S_1^1$ ($1 \leq j \leq x_1$), we proceed to find $e_{14}, e_{15}, \dots, e_{1x_1}$ in the same
 553 way, and obtain graphs $G_{1j} = G - \{e_{11}, e_{12}, \dots, e_{1(j-1)}\}$ ($1 \leq j \leq x_1$). Let
 554 $M_1 = \{e_{11}, e_{12}, \dots, e_{1x_1}\}$ and $G_1 = G - M_1$. Thus the tree T_1 induced by the
 555 edges in $\{w_1 u_{x_2+1}, w_1 u_{x_2+2}, \dots, w_1 u_k\} \cup \{e_{11}, e_{12}, \dots, e_{1x_1}\}$ is our desired tree.

556 Let us now prove the following claim.

557 **Claim 1.** $\delta(G_1[S]) \geq \frac{k-2}{2}$.

558 *Proof of Claim 1.* Assume, to the contrary, that $\delta(G_1[S]) \leq \frac{k-4}{2}$. Then there
 559 exists a vertex $u_p \in S$ such that $d_{G_1[S]}(u_p) \leq \frac{k-4}{2}$. If $u_p \in S_2^1$, then by our
 560 procedure $d_G(u_p) = d_{G_1[S]}(u_p) + 1 \leq \frac{k-2}{2}$, which implies that $d_{M \cap K_n[S]}(u_p) \geq$
 561 $k-1 - \frac{k-2}{2} = \frac{k}{2}$. Since $w_1 u_p \in M$, it follows that $d_{K_n[M]}(u_p) \geq d_{M \cap K_n[S]}(u_p) + 1 \geq$
 562 $\frac{k+2}{2}$, which contradicts $\Delta(K_n[M]) \leq \frac{k}{2}$. Let us now assume $u_p \in S_2^1$. By the above
 563 procedure, there exists a vertex $u_q \in S_1^1$ such that when we select the edge $e_{1j} =$
 564 $u_p u_q$ ($1 \leq j \leq x_1$) from $G_{1(j-1)}[S]$ the degree of u_p in $G_{1j}[S]$ is equal to $\frac{k-4}{2}$. Thus,
 565 $d_{G_{1j}[S]}(u_p) = \frac{k-4}{2}$ and $d_{G_{1(j-1)}[S]}(u_p) = \frac{k-2}{2}$. From our procedure, $|E_G[u_q, S_2^1]| =$
 566 $|E_{G_{1(j-1)}}[u_q, S_2^1]|$. Without loss of generality, let $|E_G[u_q, S_2^1]| = t$ and $u_q u_j \in E(G)$
 567 for $x_1 + 1 \leq j \leq x_1 + t$; see Figure 3 (a). Thus $u_p \in \{u_{x_1+1}, u_{x_1+2}, \dots, u_{x_1+t}\}$,
 568 and $u_q u_j \in M$ for $x_1 + t + 1 \leq j \leq k$. Because $|E_G[u_j, S_2^1]| \geq 1$ for each $u_j \in$
 569 S_1^1 ($1 \leq j \leq x_1$), we have $t \geq 1$. Since $|M| = k-1$ and $u_j w_1 \in M$ for $1 \leq j \leq x_1$,
 570 it follows that $1 \leq t \leq k-2$. Since $d_{G_{1(j-1)}[S]}(u_p) = \frac{k-2}{2}$, by our procedure
 571 $d_{G_{1(j-1)}[S]}(u_j) \leq \frac{k-2}{2}$ for each $u_j \in S_2^1$ ($x_1 + 1 \leq j \leq x_1 + t$). Assume, to the
 572 contrary, that there is a vertex u_s ($x_1 + 1 \leq s \leq x_1 + t$) such that $d_{G_{1(j-1)}[S]}(u_s) \geq$
 573 $\frac{k-2}{2}$. Then we should have selected the edge $u_q u_s$ instead of $e_{1j} = u_q u_p$ by our
 574 procedure, a contradiction. We conclude that $d_{G_{1(j-1)}[S]}(u_r) \leq \frac{k-2}{2}$ for each
 575 $u_r \in S_1^1$ ($x_1 + 1 \leq r \leq x_1 + t$). Clearly, there are at least $k-1 - \frac{k-2}{2} = \frac{k}{2}$ edges
 576 incident to each u_r ($x_1 + 1 \leq r \leq x_1 + t$) belonging to $M \cup \{e_{11}, e_{12}, \dots, e_{1(j-1)}\}$.

577 Since $j \leq x_1$ and $u_q u_j \in M$ for $x_i + t + 1 \leq j \leq k$, we have

$$\begin{aligned} & |E_{K_n[M]}[u_q, S_2^1]| + \sum_{j=1}^t d_{K_n[M]}(u_j) \\ & \geq k - x_1 - t + \frac{k}{2}t - (j - 1) - \binom{t}{2} \\ & = k + \frac{(k-2)}{2}t - x_1 - j + 1 - \binom{t}{2} \end{aligned}$$

578 and hence

$$\begin{aligned} |M| & \geq |M \cap (E_{K_n}[w_1, S])| + \sum_{j=1}^t d_{K_n[M]}(u_j) + |E_{K_n[M]}[u_q, S_1^1]| \\ & \geq x_1 + \left(k + \frac{(k-2)}{2}t - x_1 - j + 1 \right) - \binom{t}{2} \\ & = -\frac{t^2}{2} + \frac{t}{2} + \frac{(k-2)}{2}t + k - j + 1 \\ & = -\frac{t^2}{2} + \frac{(k-1)}{2}t + k - j + 1 \\ & = -\frac{1}{2} \left(t - \frac{k-1}{2} \right)^2 + \frac{(k-1)^2}{8} + k - j + 1 \\ & \geq \frac{k}{2} - 1 + k - j + 1 && \text{(since } 1 \leq t \leq k - 2) \\ & = \frac{k}{2} + k - j \\ & \geq k, && \left(\text{since } j \leq x_1 \text{ and } x_1 \leq \frac{k}{2} \right) \end{aligned}$$

contradicting $|M| = k - 1$.




Figure 3. Graphs for Lemma 2.8.

579

580 By Claim 1, we have $\delta(G_1[S]) \geq \frac{k-2}{2}$. Recall that there exists at most one
 581 vertex in $K_n[M]$ such that its degree is $\frac{k}{2}$, and $x_{n-k} \leq x_{n-k-1} \leq \cdots \leq x_2 \leq$
 582 $x_1 \leq \frac{k}{2}$. Then $x_i \leq \frac{k-2}{2}$ for $2 \leq i \leq n-k$. Now we continue to introduce our
 583 procedure.

584

For $w_2 \in \bar{S}$, without loss of generality, let $S = S_1^2 \cup S_2^2$ and $S_1^2 = \{u_1, u_2, \dots,$
 585 $u_{x_2}\}$ such that $u_j w_2 \in M$ for $1 \leq j \leq x_2$. Let $S_2^2 = S - S_1^2 = \{u_{x_2+1}, u_{x_2+2}, \dots, u_k\}$.
 586 Then $u_j w_2 \in E(G)$ for $x_2+1 \leq j \leq k$. Clearly, the tree T'_2 induced by the edges in
 587 $\{w_2 u_{x_2+1}, w_2 u_{x_2+2}, \dots, w_2 u_k\}$ is a Steiner tree connecting S_2^2 . Our idea is to seek
 588 for x_2 edges in $E_{G_1}[S_1^2, S_2^2]$ and add them to T'_2 to form a Steiner tree connecting
 589 S . For each $u_j \in S_1^2$ ($1 \leq j \leq x_2$), we claim that $|E_{G_1}[u_j, S_2^2]| \geq 1$. Otherwise, let
 590 $|E_{G_1}[u_j, S_2^2]| = 0$. Recall that $|M_1| = x_1$. Then there exist $k-x_2$ edges between u_j
 591 and S_2^2 belonging to $M \cup M_1$, and hence $|E_{K_n[M]}[u_j, S_2^2]| \geq k-x_2-x_1$. Therefore,
 592 $|M| \geq |E_{K_n[M]}[u_j, S_2^2]| + d_{K_n[M]}(w_1) + d_{K_n[M]}(w_2) \geq (k-x_2-x_1) + x_1 + x_2 = k$,
 593 which contradicts $|M| = k-1$. Choose the vertex with the smallest subscript
 594 among all the vertices of S_1^2 having maximum degree in $G_1[S]$, say u'_1 . Then
 595 we select the vertex adjacent to u'_1 with the smallest subscript among all the
 596 vertices of S_2^2 having maximum degree in $G_1[S]$, say u''_1 . Let $e_{21} = u'_1 u''_1$. Con-
 597 sider the graph $G_{21} = G_1 - e_{21}$, and choose the one with the smallest sub-
 598 script among all the vertices of $S_1^2 - u'_1$ having maximum degree in $G_{21}[S]$, say
 599 u'_2 . Then we select the vertex adjacent to u'_2 with the smallest subscript a-
 600 mong all the vertices of S_2^2 having maximum degree in $G_{21}[S]$, say u''_2 . Set
 601 $e_{22} = u'_2 u''_2$. Consider the graph $G_{22} = G_{21} - e_{22} = G_1 - \{e_{21}, e_{22}\}$. For
 602 each $u_j \in S_1^2$ ($1 \leq j \leq x_2$), we proceed to find $e_{23}, e_{24}, \dots, e_{2x_2}$ in the same
 603 way, and get graphs $G_{2j} = G_1 - \{e_{21}, e_{22}, \dots, e_{2(j-1)}\}$ ($1 \leq j \leq x_2$). Let
 604 $M_2 = \{e_{21}, e_{22}, \dots, e_{2x_2}\}$ and $G_2 = G_1 - M_1$. Thus the tree T_2 induced by
 605 the edges in $\{w_2 u_{x_2+1}, w_2 u_{x_2+2}, \dots, w_2 u_k\} \cup \{e_{21}, e_{22}, \dots, e_{2x_2}\}$ is our desired
 606 tree. Furthermore, T_2 and T_1 are two internally disjoint S -Steiner trees.

607

For $w_i \in \bar{S}$, without loss of generality, let $S = S_1^i \cup S_2^i$ and $S_1^i = \{u_1, u_2, \dots,$
 608 $u_{x_i}\}$ such that $u_j w_i \in M$ for $1 \leq j \leq x_i$. Set $S_2^i = S - S_1^i = \{u_{x_i+1}, u_{x_i+2}, \dots, u_k\}$.
 609 Then $u_j w_i \in E(G)$ for $x_i+1 \leq j \leq k$. One can see that the tree T'_i induced by the
 610 edges in $\{w_i u_{x_i+1}, w_i u_{x_i+2}, \dots, w_i u_k\}$ is a Steiner tree connecting S_2^i . Our idea
 611 is to seek for x_i edges in $E_{G_{i-1}}[S_1^i, S_2^i]$ and add them to T'_i to form a Steiner tree
 612 connecting S . For each $u_j \in S_1^i$ ($1 \leq j \leq x_i$), we claim that $|E_{G_{i-1}}[u_j, S_2^i]| \geq 1$.
 613 Otherwise, let $|E_{G_{i-1}}[u_j, S_2^i]| = 0$. Recall that $|M_j| = x_j$ ($1 \leq j \leq i$). Then
 614 there are $k-x_i$ edges between u_j and S_2^i belonging to $M \cup (\bigcup_{j=1}^{i-1} M_j)$, and
 615 hence $|E_{K_n[M]}[u_j, S_2^i]| \geq k-x_i - \sum_{j=1}^{i-1} x_j$. Therefore, $|M| \geq |E_{K_n[M]}[u_j, S_2^i]| +$
 616 $\sum_{j=1}^i |M \cap (K_n[w_j, S])| \geq k-x_i - \sum_{j=1}^{i-1} x_j + \sum_{j=1}^i x_j = k$, contradicting $|M| =$
 617 $k-1$. Choose the vertex with the smallest subscript among all the vertices of S_1^i
 618 having maximum degree in $G_{i-1}[S]$, say u'_1 . Then we select the vertex adjacent
 619 to u'_1 with the smallest subscript among all the vertices of S_2^i having maximum

620 degree in $G_{i-1}[S]$, say u'_1 . Let $e_{i1} = u'_1 u''_1$. Consider the graph $G_{i1} = G_{i-1} - e_{i1}$,
 621 choose the vertex with the smallest subscript among all the vertices of $S_1^i - u'_1$
 622 having maximum degree in $G_{i1}[S]$, say u'_2 . Then we select the vertex adjacent
 623 to u'_2 with the smallest subscript among all the vertices of S_2^i having maximum
 624 degree in $G_{i1}[S]$, say u''_2 . Set $e_{i2} = u'_2 u''_2$. Consider the graph $G_{i2} = G_{i1} - e_{i2} =$
 625 $G_{i-1} - \{e_{i1}, e_{i2}\}$. For each $u_j \in S_1^i$ ($1 \leq j \leq x_i$), we proceed to find $e_{i3}, e_{i4}, \dots, e_{ix_i}$
 626 in the same way, and get graphs $G_{ij} = G_{i-1} - \{e_{i1}, e_{i2}, \dots, e_{i(j-1)}\}$ ($1 \leq j \leq x_i$).
 627 Let $M_i = \{e_{i1}, e_{i2}, \dots, e_{ix_i}\}$ and $G_i = G_{i-1} - M_i$. Thus the tree T_i induced by
 628 the edges in $\{w_i u_{x_2+1}, w_i u_{x_2+2}, \dots, w_i u_k\} \cup \{e_{i1}, e_{i2}, \dots, e_{ix_i}\}$ is our desired tree.
 629 Furthermore, T_1, T_2, \dots, T_i are pairwise internally disjoint S -Steiner trees.

630 We continue this procedure until we obtain $n - k$ pairwise internally disjoint
 631 trees T_1, T_2, \dots, T_{n-k} . Note that if there exists some x_j such that $x_j = 0$ then
 632 $x_{j+1} = x_{j+2} = \dots = x_{n-k} = 0$ since $x_1 \geq x_2 \geq \dots \geq x_{n-k}$. Then the trees T_i
 633 induced by the edges in $\{w_i u_1, w_i u_2, \dots, w_i u_k\}$ ($j \leq i \leq n - k$) is our desired tree.
 634 From the above procedure, the resulting graph must be $G_{n-k} = G - \bigcup_{i=1}^{n-k} M_i$.
 635 Let us show the following claim.

636 **Claim 2.** $\delta(G_{n-k}[S]) \geq \frac{k-2}{2}$.

637 **Proof of Claim 2.** Assume, to the contrary, that $\delta(G_{n-k}[S]) \leq \frac{k-4}{2}$, namely,
 638 there exists a vertex $u_p \in S$ such that $d_{G_{n-k}[S]}(u_p) \leq \frac{k-4}{2}$. Since $\delta(G[S]) \geq \frac{k-2}{2}$,
 639 by our procedure there exists an edge e_{ij} in $G_{i(j-1)}$ incident to the vertex u_p such
 640 that when we pick up this edge, $d_{G_{ij}[S]}(u_p) = \frac{k-4}{2}$ but $d_{G_{i(j-1)}[S]}(u_p) = \frac{k-2}{2}$.

641 First, we consider the case $u_p \in S_2^i$. Then there exists a vertex $u_q \in S_1^i$
 642 such that when we select the edge $e_{ij} = u_p u_q$ from $G_{i(j-1)}[S]$ the degree of
 643 u_p in $G_{ij}[S]$ is equal to $\frac{k-4}{2}$. Thus, $d_{G_{ij}[S]}(u_p) = \frac{k-4}{2}$ and $d_{G_{i(j-1)}[S]}(u_p) =$
 644 $\frac{k-2}{2}$. From our procedure, $|E_{G_{i-1}}[u_q, S_2^i]| = |E_{G_{i(j-1)}}[u_q, S_2^i]|$. Without loss of
 645 generality, let $|E_{G_{i-1}}[u_q, S_2^i]| = t$ and $u_q u_j \in E(G_{i-1})$ for $x_i + 1 \leq j \leq x_i + t$; see
 646 Figure 3 (b). Thus $u_p \in \{u_{x_i+1}, u_{x_i+2}, \dots, u_{x_i+t}\}$, and $u_q u_j \in M \cup (\bigcup_{r=1}^{i-1} M_r)$ for
 647 $x_i + t + 1 \leq j \leq k$. Since $x_i \leq \frac{k-2}{2}$ ($2 \leq i \leq n - k$), it follows that $|S_1^i| \leq \frac{k-2}{2}$.
 648 From this together with $\delta(G_{i-1}[S]) \geq \frac{k-2}{2}$, we have $|E_{G_{i-1}}[u_q, S_1^i]| \geq 1$, that is,
 649 $t \geq 1$. Since $d_{G_{i(j-1)}[S]}(u_p) = \frac{k-2}{2}$, by our procedure $d_{G_{i(j-1)}[S]}(u_j) \leq \frac{k-2}{2}$ for each
 650 $u_j \in S_2^i$ ($x_i + 1 \leq j \leq x_i + t$). Assume, to the contrary, that there exists a vertex
 651 u_s ($x_i + 1 \leq s \leq x_i + t$) such that $d_{G_{i(j-1)}[S]}(u_s) \geq \frac{k-2}{2}$. Then we should have
 652 selected the edge $u_q u_s$ instead of $e_{ij} = u_q u_p$ by our procedure, a contradiction.
 653 We conclude that $d_{G_{i(j-1)}[S]}(u_r) \leq \frac{k-2}{2}$ for each $u_r \in S_2^i$ ($x_i + 1 \leq r \leq x_i + t$).
 654 Clearly, there are at least $k - 1 - \frac{k-2}{2} = \frac{k}{2}$ edges incident to each u_r ($x_i + 1 \leq$
 655 $r \leq x_i + t$) belonging to $M \cup (\bigcup_{j=1}^{i-1} M_j) \cup \{e_{i1}, e_{i2}, \dots, e_{i(j-1)}\}$. Since $j \leq x_i$ and

656 $u_q u_j \in M \cup (\bigcup_{r=1}^{i-1} M_r)$ for $x_i + t + 1 \leq j \leq k$, we have

$$\begin{aligned}
& |E_{K_n[M]}[u_q, S_2^i]| + \sum_{j=1}^t d_{K_n[M]}(u_j) \\
\geq & k - x_i - t + \frac{k}{2}t - \sum_{j=1}^{i-1} x_j - (j-1) - \binom{t}{2} \\
\geq & k + \frac{(k-2)}{2}t - \sum_{j=1}^i x_j - x_i + 1 - \binom{t}{2} \quad (\text{since } j \leq x_i) \\
= & -\frac{t^2}{2} + \frac{(k-1)}{2}t + k - \sum_{j=1}^i x_j - x_i + 1 \\
= & -\frac{1}{2} \left(t - \frac{k-1}{2} \right)^2 + \frac{(k-1)^2}{8} + k - \sum_{j=1}^i x_j - x_i + 1
\end{aligned}$$

657 and hence

$$\begin{aligned}
|M| & \geq \sum_{j=1}^i |M \cap (E_{K_n}[w_j, S])| + \sum_{j=1}^t d_{K_n[M]}(u_j) + |E_{K_n[M]}[u_q, S_2^i]| \\
& \geq \sum_{j=1}^i x_j - \frac{1}{2} \left(t - \frac{k-1}{2} \right)^2 + \frac{(k-1)^2}{8} + k - \sum_{j=1}^i x_j - x_i + 1 \\
& = -\frac{1}{2} \left(t - \frac{k-1}{2} \right)^2 + \frac{(k-1)^2}{8} + k - x_i + 1 \\
& \geq \frac{k}{2} - 1 + k - x_i + 1 \quad (\text{since } 1 \leq t \leq k-2) \\
& \geq \frac{k}{2} + k - x_i \\
& \geq k + 1, \quad \left(\text{since } x_i \leq \frac{k-2}{2} \right)
\end{aligned}$$

658 which contradicts $|M| = k - 1$.

659 Next, assume $u_p \in S_1^i$. Recall that $d_{G_{ij}[S]}(u_p) = \frac{k-4}{2}$. Since $u_p \in S_1^i$, it
660 follows that $d_{G_{i-1}[S]}(u_p) = \frac{k-2}{2}$. If $u_p \in \bigcap_{j=1}^i S_1^j$, namely, $u_p w_j \in M$ ($1 \leq j \leq i$),
661 then by our procedure $d_{G[S]}(u_p) = \frac{k-2}{2} + i - 1$ and hence $d_{K_n[S] \cap M}(u_p) = k - 1 -$
662 $(\frac{k-2}{2} + i - 1) = \frac{k}{2} - i + 1$. Since $u_p w_j \in M$ for each $w_j \in \bar{S}$ ($1 \leq j \leq i$), we have
663 $d_{K_n[M]}(u_p) = d_{K_n[S] \cap M}(u_p) + d_{K_n[S, \bar{S}] \cap M}(u_p) \geq (\frac{k}{2} - i + 1) + i = \frac{k+2}{2}$, contradicting
664 $\Delta(K_n[M]) \leq \frac{k}{2}$. Combining this with $u_p \in S_1^i$, we have $u_p \notin \bigcap_{j=1}^{i-1} S_1^j$ and we

665 can assume that there exists an integer i' ($i' \leq i - 1$) satisfying the following
 666 conditions:

- 667 • $u_p \in S_2^{i'}$ and $d_{G_{i'}[S]}(u_p) < d_{G_{i'-1}[S]}(u_p)$;
- 668 • if u_p belongs to some S_2^j ($i' + 1 \leq j \leq i$) then $d_{G_j[S]}(u_p) = d_{G_{j-1}[S]}(u_p)$.

669 The above two conditions can be restated as follows:

- 670 • $u_p w_{i'} \in E(G)$ and $d_{G_{i'}[S]}(u_p) < d_{G_{i'-1}[S]}(u_p)$;
- 671 • if $u_p w_j \in E(G)$ ($i' + 1 \leq j \leq i$) then $d_{G_j[S]}(u_p) = d_{G_{j-1}[S]}(u_p)$.

672 In fact, we can find the integer i' such that $u_p w_{i'} \in E(G)$ and $d_{G_{i'}[S]}(u_p) <$
 673 $d_{G_{i'-1}[S]}(u_p)$. Assume, to the contrary, that for each w_j ($1 \leq j \leq i$), $u_p w_j \in M$,
 674 or $u_p w_j \in E(G)$ but $d_{G_j[S]}(u_p) = d_{G_{j-1}[S]}(u_p)$. Let i_1 ($i_1 \leq i$) be the number
 675 of vertices nonadjacent to $u_p \in S$ in $\{w_1, w_2, \dots, w_{i-1}\} \subseteq \bar{S}$. Without loss of
 676 generality, let $w_j u_p \in M$ ($1 \leq j \leq i_1$). Recall that $d_{G_{ij}[S]}(u_p) = \frac{k-4}{2}$. Thus
 677 $d_{G[S]}(u_p) = \frac{k-4}{2} + i_1$ and hence $d_{K_n[S] \cap M}(u_p) \geq k - 1 - (\frac{k-4}{2} + i_1) = \frac{k+2}{2} -$
 678 i_1 . Since $w_j u_p \in M$ ($1 \leq j \leq i_1$), it follows that $d_{K_n[S, \bar{S}] \cap M}(u_p) \geq i_1$, which
 679 results in $d_{K_n[M]}(u_p) = d_{K_n[S] \cap M}(u_p) + d_{K_n[S, \bar{S}] \cap M}(u_p) \geq (\frac{k+2}{2} - i_1) + i_1 = \frac{k+2}{2}$,
 680 contradicting $\Delta(K_n[M]) \leq \frac{k}{2}$.

681 Now we turn our attention to $u_p \in S_2^{i'}$. Without loss of generality, let
 682 $u_p w_j \in M$ ($j \in \{j_1, j_2, \dots, j_{i_1}\}$), namely, $u_p \in S_1^{j_1} \cap S_1^{j_2} \cap \dots \cap S_1^{j_{i_1}}$, where
 683 $j_1, j_2, \dots, j_{i_1} \in \{i'+1, i'+2, \dots, i\}$. Then $u_p w_j \in E(G)$ ($j \in \{i'+1, i'+2, \dots, i\} -$
 684 $\{j_1, j_2, \dots, j_{i_1}\}$). Clearly, $i_1 \leq i - i'$. Recall that $u_p \in S_1^i$ and $d_{G_{ij}[S]}(u_p) = \frac{k-4}{2}$.
 685 Thus $d_{G_{i'}[S]}(u_p) = \frac{k-4}{2} + i_1$. By our procedure, there exists a vertex $u_q \in S_1^{i'}$
 686 such that when we select the edge $e_{i'j} = u_p u_q$ from $G_{i'(j-1)}[S]$ the degree of u_p in
 687 $G_{i'j}[S]$ is equal to $\frac{k-4}{2} + i_1$, that is, $d_{G_{i'j}[S]}(u_p) = \frac{k-4}{2} + i_1$ and $d_{G_{i'(j-1)}[S]}(u_p) =$
 688 $\frac{k-2}{2} + i_1$. From our procedure, $|E_{G_{i'-1}}[u_q, S_2^{i'}]| = |E_{G_{i'(j-1)}}[u_q, S_2^{i'}]|$. Without
 689 loss of generality, let $|E_{G_{i'-1}}[u_q, S_2^{i'}]| = t$ and $u_q u_j \in E(G_{i'-1})$ for $x_{i'} + 1 \leq$
 690 $j \leq x_{i'} + t$; see Figure 3 (c). Thus $u_p \in \{u_{x_{i'}+1}, u_{x_{i'}+2}, \dots, u_{x_{i'}+t}\}$, and $u_q u_j \in$
 691 $M \cup (\bigcup_{r=1}^{i'-1} M_r)$ for $x_{i'} + t + 1 \leq j \leq k$. Since $x_j \leq \frac{k-2}{2}$ ($2 \leq j \leq n - k$), it
 692 follows that $|S_1^{i'}| \leq \frac{k-2}{2}$. From this together with $\delta(G_{i'-1}[S]) \geq \frac{k-2}{2}$, we have
 693 $|E_{G_{i'-1}}[u_q, S_1^{i'}]| \geq 1$, that is, $t \geq 1$. Since $d_{G_{i'(j-1)}[S]}(u_p) = \frac{k-2}{2} + i_1$, by our
 694 procedure $d_{G_{i'(j-1)}[S]}(u_j) \leq \frac{k-2}{2} + i_1$ for each $u_j \in S_2^{i'}$ ($x_{i'} + 1 \leq j \leq x_{i'} + t$).
 695 Assume, to the contrary, that there is a vertex u_s ($x_{i'} + 1 \leq s \leq x_{i'} + t$) such
 696 that $d_{G_{i'(j-1)}[S]}(u_s) \geq \frac{k-2}{2} + i_1 + 1$. Then we should have selected the edge
 697 $u_q u_s$ instead of $e_{i'j} = u_p u_q$ by our procedure, a contradiction. We conclude
 698 that $d_{G_{i'(j-1)}[S]}(u_r) \leq \frac{k-2}{2} + i_1$ for each $u_r \in S_2^{i'}$ ($x_{i'} + 1 \leq r \leq x_{i'} + t$).
 699 Clearly, there are at least $k - 1 - (\frac{k-2}{2} + i_1) = \frac{k}{2} - i_1$ edges incident to each
 700 u_r ($x_{i'} + 1 \leq r \leq x_{i'} + t$) belonging to $M \cup (\bigcup_{j=1}^{i'-1} M_j) \cup \{e_{i'1}, e_{i'2}, \dots, e_{i'(j-1)}\}$.

701 Since $j \leq x_{i'}$ and $u_q u_j \in M \cup (\bigcup_{r=1}^{i'-1} M_r)$ for $x_{i'} + t + 1 \leq j \leq k$, we have

$$\begin{aligned}
& |E_{K_n[M]}[u_q, S_2^{i'}]| + \sum_{j=1}^t d_{K_n[M]}(u_j) \\
\geq & k - x_{i'} - t + \left(\frac{k}{2} - i_1\right)t - \sum_{j=1}^{i'-1} x_j - (j-1) - \binom{t}{2} \\
\geq & k - \sum_{j=1}^{i'} x_j + \left(\frac{k-2}{2} - i_1\right)t - x_{i'} + 1 - \frac{t(t-1)}{2} \quad (\text{since } j \leq x_{i'}) \\
= & -\frac{t^2}{2} + \frac{t}{2} + k - \sum_{j=1}^{i'} x_j + \left(\frac{k-2}{2} - i + i'\right)t - x_{i'} + 1 \quad (\text{since } i_1 \leq i - i') \\
= & -\frac{t^2}{2} + \left(\frac{k-1}{2} - i + i'\right)t + k - \sum_{j=1}^{i'} x_j - x_{i'} + 1 \\
= & -\frac{1}{2}(t^2 - (k-1-2i+2i')t) + k - \sum_{j=1}^{i'} x_j - x_{i'} + 1 \\
= & -\frac{1}{2}\left(t - \frac{k-1-2i+2i'}{2}\right)^2 + \frac{(k-1-2i+2i')^2}{8} + k - \sum_{j=1}^{i'} x_j - x_{i'} + 1
\end{aligned}$$

702 and hence

$$\begin{aligned}
& |M| \\
\geq & \sum_{j=1}^i |M \cap (E_{K_n}[w_j, S])| + \sum_{j=1}^p d_{K_n[M]}(u_j) + |E_{K_n[M]}[u_q, S_2^i]| \\
\geq & \sum_{j=1}^i x_j - \frac{1}{2}\left(t - \frac{k-1-2i+2i'}{2}\right)^2 + \frac{(k-1-2i+2i')^2}{8} + k - \sum_{j=1}^{i'} x_j - x_{i'} \\
& + 1 \\
= & -\frac{1}{2}\left(t - \frac{k-1-2i+2i'}{2}\right)^2 + \frac{(k-1-2i+2i')^2}{8} + k + \sum_{j=i'+1}^i x_j - x_{i'} + 1 \\
\geq & \frac{k}{2} - 1 - i + i' + k + \sum_{j=i'+1}^i x_j - x_{i'} + 1 \quad (\text{since } 1 \leq t \leq k-2 \text{ and} \\
& k-1-2i+2i' \leq k-2) \\
\geq & k, \quad \left(\text{since } x_{i'} \leq \frac{k-2}{2} \text{ and } x_j \geq 1 \text{ for } i'+1 \leq j \leq i\right)
\end{aligned}$$

703 contradicting $|M| = k - 1$. This completes the proof of Claim 2.

704 From our procedure, we get $n - k$ internally disjoint Steiner trees connecting
 705 S in G , say T_1, T_2, \dots, T_{n-k} . Recall that $G_{n-k} = G - (\bigcup_{i=1}^{n-k} M_i)$. We can
 706 regard $G_{n-k}[S] = G[S] - (\bigcup_{i=1}^{n-k} M_i)$ as a graph obtained from the complete
 707 graph K_k by deleting $|M'| + \sum_{i=1}^{n-k} |M_i|$ edges. Since $|M'| + \sum_{i=1}^{n-k} |M_i| + |M''| =$
 708 $m_1 + \sum_{i=1}^{n-k} x_i + m_2 = k - 1$, we have $1 \leq \sum_{i=1}^{n-k} |M_i| + m_1 \leq k - 1$. By Claim 2,
 709 $\delta(G_{n-k}[S]) \geq \frac{k-2}{2}$ and hence $2 \leq \Delta(G_{n-k}[S]) \leq \frac{k}{2}$. From Lemma 2.6, there exist
 710 $\frac{k-2}{2}$ edge-disjoint spanning trees connecting S in $G_{n-k}[S]$. These trees together
 711 with T_1, T_2, \dots, T_{n-k} are $n - \frac{k}{2} - 1$ internally disjoint Steiner trees connecting S in
 712 G . Thus, $\kappa(S) \geq n - \frac{k}{2} - 1$. From the arbitrariness of S , we have $\kappa_k(G) \geq n - \frac{k}{2} - 1$,
 713 as desired. \blacksquare

714 We are now in a position to prove our main results.

715 **Proof of Theorem 1.8.** Assume that $\kappa_k(G) = n - \frac{k}{2} - 1$. Since G of order
 716 n is connected, we can regard G as a graph obtained from the complete graph
 717 K_n by deleting some edges. From Lemma 1.7, it follows that $|M| \geq 1$ and hence
 718 $\Delta(K_n[M]) \geq 1$. If $G = K_n - M$ where $M \subseteq E(K_n)$ such that $\Delta(K_n[M]) \geq$
 719 $\frac{k}{2} + 1$, then $\kappa_k(G) \leq \lambda_k(G) < n - \frac{k}{2} - 1$ by Observation 1.2 and Corollary 2.2, a
 720 contradiction. So $1 \leq \Delta(K_n[M]) \leq \frac{k}{2}$. If $2 \leq \Delta(K_n[M]) \leq \frac{k}{2}$ and $|M| \geq k$, then
 721 $\kappa_k(G) \leq \lambda_k(G) < n - \frac{k}{2} - 1$ by Observation 1.2 and Lemma 2.4, a contradiction.
 722 Therefore, $1 \leq |M| \leq k - 1$. If $\Delta(K_n[M]) = 1$, then $1 \leq |M| \leq k - 1$ by Lemma
 723 2.5. We conclude that $1 \leq \Delta(K_n[M]) \leq \frac{k}{2}$ and $1 \leq |M| \leq k - 1$, as desired.

724 Conversely, let $G = K_n - M$ where $M \subseteq E(K_n)$ such that $1 \leq \Delta(K_n[M]) \leq \frac{k}{2}$
 725 and $1 \leq |M| \leq k - 1$. In fact, we only need to show that $\kappa_k(G) \geq n - \frac{k}{2} - 1$ for
 726 $\Delta(K_n[M]) = 1$ and $|M| = k - 1$, or $2 \leq \Delta(K_n[M]) \leq \frac{k}{2}$ and $|M| = k - 1$. The
 727 results follow by (1) of Lemma 2.7 and Lemma 2.8. \square

728 **Proof of Theorem 1.9.** If G is a connected graph satisfying condition (2), then
 729 $\kappa_k(G) = n - \frac{k}{2} - 1$ by Theorem 1.8. From Observation 1.2, $\lambda_k(G) \geq \kappa_k(G) =$
 730 $n - \frac{k}{2} - 1$. From this together with Lemma 1.7, we have $\lambda_k(G) = n - \frac{k}{2} - 1$.
 731 Assume that G is a connected graph satisfying condition (1). We only need to
 732 show that $\lambda_k(G) = n - \frac{k}{2} - 1$ for $|M| = \lfloor \frac{n}{2} \rfloor$. The result follows by (2) of Lemma
 733 2.7 and Lemma 1.7.

734 Conversely, assume that $\lambda_k(G) = n - \frac{k}{2} - 1$. Since G of order n is connected,
 735 we can consider G as a graph obtained from a complete graph K_n by deleting some
 736 edges. From Corollary 2.2, $G = K_n - M$ such that $\Delta(K_n[M]) \leq \frac{k}{2}$, where $M \subseteq$
 737 $E(K_n)$. Combining this with Lemma 1.7, we have $|M| \geq 1$ and $\Delta(K_n[M]) \geq 1$.
 738 So $1 \leq \Delta(K_n[M]) \leq \frac{k}{2}$. It is clear that if $\Delta(K_n[M]) = 1$ then $1 \leq |M| \leq \lfloor \frac{n}{2} \rfloor$. If
 739 $2 \leq \Delta(K_n[M]) \leq \frac{k}{2}$, then $1 \leq |M| \leq k - 1$ by Lemma 2.4. So (1) or (2) holds. \square

740 **Remark 3.** As we know, $\lambda(G) = n - 2$ if and only if $G = K_n - M$ such that
 741 $\Delta(K_n[M]) = 1$ and $1 \leq |M| \leq \lfloor \frac{n}{2} \rfloor$, where $M \subseteq E(K_n)$. So we can restate the
 742 above conclusion as follows: $\lambda_2(G) = n - 2$ if and only if $G = K_n - M$ such
 743 that $\Delta(K_n[M]) = 1$ and $1 \leq |M| \leq \lfloor \frac{n}{2} \rfloor$, where $M \subseteq E(K_n)$. This means that
 744 $4 \leq k \leq n$ in Theorem 1.9 can be replaced by $2 \leq k \leq n$.

745 **Acknowledgements:** The authors are very grateful to the referees for valu-
 746 able comments and suggestions, which helped to improve the presentation of the
 747 paper. The author X. Li was supported by NSFC Nos. 11371205 and 11531011,
 748 and PCSIRT, and the author Y. Mao was supported by NSFC Nos. 11551001,
 749 11161037, 11461054 and SFQP No. 2014-ZJ-907.

750 REFERENCES

- 751 [1] L.W. Beineke, R.J. Wilson, *Topics in Structural Graph Theory*, Cambridge
 752 University Press, 2013.
- 753 [2] F.T. Boesch, S. Chen, *A generalization of line connectivity and optimally*
 754 *invulnerable graphs*, SIAM J. Appl. Math. 34(1978), 657–665.
- 755 [3] J.A. Bondy, U.S.R. Murty, *Graph Theory*, GTM 244, Springer, 2008.
- 756 [4] G. Chartrand, S.F. Kapoor, L. Lesniak, D.R. Lick, *Generalized connectivity*
 757 *in graphs*, Bull. Bombay Math. Colloq. 2(1984), 1–6.
- 758 [5] G. Chartrand, F. Okamoto, P. Zhang, *Rainbow trees in graphs and general-*
 759 *ized connectivity*, Networks 55(4)(2010), 360–367.
- 760 [6] D.P. Day, O.R. Oellermann, H.C. Swart, *The ℓ -connectivity function of*
 761 *trees and complete multipartite graphs*, J. Combin. Math. Combin. Comput.
 762 10(1991), 183–192.
- 763 [7] D.L. Goldsmith, *On the second-order edge-connectivity of a graph*, Congres-
 764 sus Numerantium 29(1980), 479–484.
- 765 [8] D.L. Goldsmith, *On the n th order connectivity of a graph*, Congressus Nu-
 766 merantium 32(1981), 375–382.
- 767 [9] D.L. Goldsmith, B. Manval, V. Faber, *Seperation of graphs into three com-*
 768 *ponents by removal of edges*, J. Graph Theory 4(1980), 213–218.
- 769 [10] M. Grötschel, *The Steiner tree packing problem in VLSI design*, Math. Pro-
 770 gram. 78(1997), 265–281.
- 771 [11] M. Grötschel, A. Martin, R. Weismantel, *Packing Steiner trees: A cutting*
 772 *plane algorithm and commputational results*, Math. Program. 72(1996), 125–
 773 145.

- 774 [12] M. Hager, *Pendant tree-connectivity*, J. Combin. Theory 38(1985), 179–189.
- 775 [13] M. Hager, *Path-connectivity in graphs*, Discrete Math. 59(1986), 53–59.
- 776 [14] H.R. Hind, O.R. Oellermann, *Menger-type results for three or more vertices*,
777 Congressus Numerantium 113(1996), 179–204.
- 778 [15] K. Jain, M. Mahdian, M. Salavatipour, *Packing Steiner trees*, in: Proc.
779 14th ACM-SIAM symposium on Discrete Algorithms, Baltimore, 2003,
780 266–274.
- 781 [16] M. Kriesell, *Edge-disjoint trees containing some given vertices in a graph*, J.
782 Combin. Theory, Ser.B 88(2003), 53–65.
- 783 [17] M. Kriesell, *Edge-disjoint Steiner trees in graphs without large bridges*, J.
784 Graph Theory 62(2009), 188–198.
- 785 [18] L. Lau, *An approximate max-Steiner-tree-packing min-Steiner-cut theorem*,
786 Combinatorica 27(2007), 71–90.
- 787 [19] W. Mader, *Über die maximalzahl kantendisjunkter A-wege*, Arch. Math.
788 30(1978), 325–336.
- 789 [20] W. Mader, *Über die maximalzahl kreuzungsfreier H-wege*, Arch. Math.
790 31(1978), 387–402.
- 791 [21] H. Li, X. Li, Y. Mao, *On extremal graphs with at most two internally disjoint*
792 *Steiner trees connecting any three vertices*, Bull. Malays. Math. Sci. Soc.
793 (2)37(3)(2014), 747–756.
- 794 [22] H. Li, X. Li, Y. Mao, Y. Sun, *Note on the generalized connectivity*, Ars
795 Combin. 114(2014), 193–202.
- 796 [23] H. Li, X. Li, Y. Mao, J. Yue, *Note on the spanning-tree packing number of*
797 *lexicographic product graphs*, Discrete Math. 338(5-6)(2015), 669–673.
- 798 [24] H. Li, X. Li, Y. Sun, *The generalized 3-connectivity of Cartesian product*
799 *graphs*, Discrete Math. Theor. Comput. Sci. 14(1)(2012), 43–54.
- 800 [25] S. Li, W. Li, X. Li, *The generalized connectivity of complete equipartition*
801 *3-partite graphs*, Bull. Malays. Math. Sci. Soc.(2) 37(1)(2014), 103–121.
- 802 [26] S. Li, X. Li, *Note on the hardness of generalized connectivity*, J. Combin.
803 Optimization 24(2012), 389–396.
- 804 [27] S. Li, X. Li, W. Zhou, *Sharp bounds for the generalized connectivity $\kappa_3(G)$* ,
805 Discrete Math. 310(2010), 2147–2163.

- 806 [28] X. Li, Y. Mao, *Nordhaus-Gaddum-type results for the generalized edge-*
807 *connectivity of graphs*, Discrete Appl. Math. 185(2015), 102–112.
- 808 [29] X. Li, Y. Mao, *On extremal graphs with at most ℓ internally disjoint Steiner*
809 *trees connecting any $n - 1$ vertices*, Graphs Combin. 31(6)(2015), 2231–2259.
- 810 [30] X. Li, Y. Mao, *The generalized 3-connectivity of lexicographical product*
811 *graphs*, Discrete Math. Theor. Comput. Sci. 16(1)(2014), 339–354.
- 812 [31] X. Li, Y. Mao, *The minimal size of a graph with given generalized 3-edge-*
813 *connectivity*, Ars Combin. 118(2015), 63–72.
- 814 [32] X. Li, Y. Mao, Y. Sun, *On the generalized (edge-)connectivity of graphs*,
815 Australasian J. Combin. 58(2)(2014), 304–319.
- 816 [33] C.St.J.A. Nash-Williams, *Edge-disjoint spanning trees of finite graphs*, J.
817 London Math. Soc. 36(1961), 445–450.
- 818 [34] O.R. Oellermann, *Connectivity and edge-connectivity in graphs: A survey*,
819 Congressus Numerantium 116 (1996), 231–252.
- 820 [35] O.R. Oellermann, *On the ℓ -connectivity of a graph*. Graphs and Combin.
821 3(1987), 285–299.
- 822 [36] O.R. Oellermann, *A note on the ℓ -connectivity function of a graph*, Conges-
823 sus Numerantium 60(1987), 181–188.
- 824 [37] F. Okamoto, P. Zhang, *The tree connectivity of regular complete bipartite*
825 *graphs*, J. Combin. Math. Combin. Comput. 74(2010), 279–293.
- 826 [38] N.A. Sherwani, *Algorithms for VLSI Physical Design Automation*, 3rd Edi-
827 tion, Kluwer Acad. Pub., London, 1999.
- 828 [39] W. Tutte, *On the problem of decomposing a graph into n connected factors*,
829 J. London Math. Soc. 36(1961), 221–230.
- 830 [40] D. West, H. Wu, *Packing Steiner trees and S -connectors in graphs*, J. Com-
831 bin. Theory, Ser.B 102(2012), 186–205.

832 **Appendix: An example for Case 2 of Lemma 2.8**

833 Let $k = 8$ and let $G = K_n - M$ where $M \subseteq E(K_n)$ be a connected
834 graph of order n such that $|M| = k - 1 = 7$ and $\Delta(K_n[M]) \leq \frac{k}{2} = 4$.
835 Let $S = \{u_1, u_2, \dots, u_8\}$, $\bar{S} = V(G) - S = \{w_1, w_2, \dots, w_{n-8}\}$ and

836 $M = \{w_1u_1, w_1u_2, w_1u_3, w_2u_2, w_2u_4, u_5u_6, u_6u_8\}$; see Figure 4 (a). Clear-
 837 ly, $x_1 = |E_{K_n[M]}[w_1, S]| = 3 \geq x_2 = |E_{K_n[M]}[w_2, S]| = 2 > x_i =$
 838 $|E_{K_n[M]}[w_i, S]| = 0$ ($3 \leq i \leq n - 8$).

839 For w_1 , we let $S_1^1 = \{u_1, u_2, u_3\}$ since $w_1u_1, w_1u_2, w_1u_3 \in M$. Set $S_2^1 = S -$
 840 $S_1^1 = \{u_4, u_5, u_6, u_7, u_8\}$. Clearly, $d_{G[S]}(u_1) = d_{G[S]}(u_2) = d_{G[S]}(u_3) = 7 =$
 841 $k - 1$ and hence u_1, u_2, u_3 are all the vertices of S_1^1 having maximum degree in
 842 $G[S]$. But u_1 is the one with the smallest subscript, so we choose $u'_1 = u_1$ in
 843 S_1^1 and select the vertex adjacent to u'_1 in S_2^1 and obtain $u_4, u_5, u_6, u_7, u_8 \in S_2^1$
 844 since $u'_1u_j \in E(G)$ ($j = 4, \dots, 8$). Obviously, $d_{G[S]}(u_4) = d_{G[S]}(u_7) = 7 >$
 845 $d_{G[S]}(u_5) = d_{G[S]}(u_8) = 6 > d_{G[S]}(u_6) = 5$ and hence u_4, u_7 are two vertices
 846 of S_2^1 having maximum degree in $G[S]$. Since u_4 is the one with the smallest
 847 subscript, we choose $u''_1 = u_4 \in S_2^1$ and put $e_{11} = u'_1u''_1 (= u_1u_4)$. Consider the
 848 graph $G_{11} = G - e_{11}$. Since $d_{G_{11}[S]}(u_2) = d_{G_{11}[S]}(u_3) = 7$ and the subscript of
 849 u_2 is smaller than u_3 , we let $u'_2 = u_2$ in $S_1^1 - u'_1$ and select the vertices adjacent
 850 to u'_2 in S_2^1 and obtain $u_4, u_5, u_6, u_7, u_8 \in S_2^1$ since $u'_2u_j \in E(G_{11})$ ($j =$
 851 $4, \dots, 8$). Since $d_{G_{11}[S]}(u_7) = 7 > d_{G_{11}[S]}(u_j) = 6 > d_{G_{11}[S]}(u_6) = 5$ ($j =$
 852 $4, 5, 8$), we select $u''_2 = u_7 \in S_2^1$ and get $e_{12} = u'_2u''_2 (= u_2u_7)$. Consider
 853 the graph $G_{12} = G_{11} - e_{12} = G - \{e_{11}, e_{12}\}$. There is only one vertex u_3
 854 in $S_1 - \{u'_1, u'_2\} = S_1 - \{u_1, u_2\}$. Therefore, let $u'_3 = u_3$ and select the
 855 vertices adjacent to u'_3 in S_2^1 and obtain $u_j \in S_2^1$ since $u'_3u_j \in E(G_{12})$ ($j =$
 856 $4, \dots, 8$). Since $d_{G_{12}[S]}(u_j) = 6 > d_{G_{12}[S]}(u_6) = 5$ ($i = 4, 5, 7, 8$), it follows
 857 that u_4, u_5, u_7, u_8 are all the vertices of S_2^1 having maximum degree in $G_{12}[S]$.
 858 But u_4 is the one with the smallest subscript, so we choose $u''_3 = u_4 \in S_2^1$
 859 and get $e_{13} = u'_3u''_3 (= u_3u_4)$. Since $x_1 = |E_{K_n[M]}[w_1, S]| = 3$, we terminate
 860 this procedure. Set $M_1 = \{e_{11}, e_{12}, e_{13}\}$ and $G_1 = G - M_1$. Thus the tree T_1
 861 induced by the edges in $\{w_1u_4, w_1u_5, w_1u_6, w_1u_7, w_1u_8, u_1u_4, u_2u_7, u_3u_4\}$ is
 862 our desired tree; see Figure 4 (b).

863 For w_2 , we let $S_1^2 = \{u_2, u_4\}$ since $w_2u_2, w_2u_4 \in M$. Let $S_2^2 = S - S_1^2 =$
 864 $\{u_1, u_3, u_5, u_6, u_7, u_8\}$. Since $d_{G_1[S]}(u_2) = 6 > d_{G_1[S]}(u_4) = 5$, it follows
 865 that u_2 is the vertex of S_1^2 having maximum degree in $G_1[S]$. So we choose
 866 $u'_1 = u_2$ in S_1^2 and find the vertices adjacent to $u'_1 (= u_2)$ in S_2^2 and ob-
 867 tain $u_1, u_3, u_5, u_6, u_8 \in S_2^2$ since $u'_1u_j \in E(G_{21})$ ($j = 1, 3, 5, 6, 8$). Since
 868 $d_{G_1[S]}(u_j) = 6 > d_{G_1[S]}(u_6) = 5$ ($j = 1, 3, 5, 8$) and u_1 is the vertex hav-
 869 ing maximum degree with the smallest subscript, we choose $u''_1 = u_1 \in S_2^2$.
 870 Put $e_{21} = u'_1u''_1 (= u_2u_1)$. Consider the graph $G_{21} = G_1 - e_{21}$. Clearly,
 871 $S_1 - \{u'_1\} = S_1 - \{u_2\} = \{u_4\}$, so we let $u'_2 = u_4$ and select the ver-
 872 tices adjacent to $u'_2 (= u_4)$ in S_2^2 and obtain u_5, u_6, u_7, u_8 since $u_2u_j \in$
 873 $E(G)$ ($j = 5, 6, 7, 8$). Since $d_{G_{21}[S]}(u_j) = 6 > d_{G_{21}[S]}(u_6) = 5$ ($j = 5, 7, 8$)
 874 and u_5 is the vertex with the smallest subscript, we let $u''_2 = u_5 \in S_2^2$ and
 875 get $e_{22} = u'_2u''_2 (= u_4u_5)$. Since $x_2 = |E_{K_n[M]}[w_2, S]| = 2$, we terminate
 876 this procedure. Let $M_2 = \{e_{21}, e_{22}\}$ and $G_2 = G_1 - M_2$. Then the tree T_2




Figure 4 Graphs for the appendix.

877 induced by the edges in $\{w_2u_1, w_2u_3, w_2u_5, w_2u_6, w_2u_7, w_2u_8, u_2u_1, u_4u_5\}$ is
 878 our desired tree; see Figure 4 (c). Obviously, T_2 and T_1 are two internally
 879 disjoint Steiner trees connecting S .

880 Since $x_i = |E_{K_n[M]}[w_i, S]| = 0$ for $3 \leq i \leq n - 8$, we terminate
 881 this procedure. For w_3, \dots, w_{n-8} , the trees T_i induced by the edges
 882 $\{w_iu_1, w_iu_2, \dots, w_iu_8\}$ ($3 \leq i \leq n - 8$) (see Figure 4 (d)) are our desired
 883 trees.

884 We can consider $G_2[S] = G[S] - \{M_1, M_2\}$ as a graph obtained from complete
 885 graph K_k by deleting $|M \cap K_n[S]| + |M_1| + |M_2|$ edges. Since $|M \cap K_n[S]| +$
 886 $|M_1| + |M_2| = 2 + 3 + 2 = 7 = k - 1$, it follows from Lemma ?? that there exist
 887 three edge-disjoint spanning trees connecting S in $G[S]$ (Actually, we can give
 888 three edge-disjoint spanning trees; see Figure 4 (e). For example, the trees
 889 $T'_1 = u_1u_8 \cup u_8u_4 \cup u_4u_6 \cup u_6u_3 \cup u_3u_2 \cup u_2u_5 \cup u_5u_7$, $T'_2 = u_4u_7 \cup u_7u_8 \cup u_8u_3 \cup$
 890 $u_3u_1 \cup u_1u_5 \cup u_1u_6 \cup u_6u_2$ and $T'_3 = u_2u_4 \cup u_2u_8 \cup u_8u_5 \cup u_5u_3 \cup u_3u_7 \cup u_1u_7 \cup u_7u_6$
 891 can be our desired trees). These three trees together with T_1, T_2, \dots, T_{n-8}
 892 are $n - 5 = n - \frac{k}{2} - 1$ internally disjoint Steiner trees connecting S . Thus,
 893 $\lambda(S) \geq n - 5$.